Effect of Amino Acid on Panicle Length and Fruit Set of Tree Mangoes (Cultivar Tommy Atkins) in Semi Arid of North of Minas Gerais

M.C.T. Pereira, S. Nietzsche and G.P. Mizobutsi

Agrarian Science Department, Universidade Estadual de Montes Claros – UNIMONTES, CP 91, CEP 39440-000, Janaúba, MG, Brazil, E-mail: marlon.pereira@unimontes.br

E.D. Júnior and S.M. de Souza

Students of Universidade Estadual de Montes Claros.

M.A. Mouco

Embrapa Researcher

Keywords: mango, panicle, amino acid

ABSTRACT

The control of the vegetative growth and the adequacy of the time of production are two of the most important and challenging aspects of the modern and efficient handling of orchards. Paclobutrazol (PBZ) has shown itself as a product of great value for the mango culture, therefore the continuous use, causes a compacting in the panicle of the mango tree. Twenty uniform and healthy plants were selected, after that 20 branches in each plant were labeled. Three amino acid concentrations were tested, through foliar sprays. The dose of 0,15% of amino acids represents the maximum economic efficiency in the length of panicles, with 30cm and the fruit setting was not affected by the amino acid application, specifically for product AMINON-25.

BIOGRAPHY

Marlon Cristian Toledo Pereira: A registered Agronomy engineer. His Doctor Science it is in Minas Gerais, Brazil. Actually he is teacher of State University of Montes Claros, Member of Committee of seeds and seedlings of Minas Gerais State.