

## Notas Científicas

### Temperature changes in soil covered by black oat straw

Anderson Luiz Zwirtes<sup>(1)</sup>, Dalvan José Reinert<sup>(2)</sup>, Paulo Ivonir Gubiani<sup>(2)</sup>,  
Vanderlei Rodrigues Da Silva<sup>(3)</sup>, Rodrigo Pivoto Mulazzani<sup>(2)</sup> and André Somavilla<sup>(2)</sup>

<sup>(1)</sup>Instituto Federal Santa Catarina, Campus de São Miguel do Oeste, Rua 22 de Abril, nº 2440, São Luiz, CEP 89900-000, São Miguel do Oeste, SC, Brazil. E-mail: andersonzwirtes@yahoo.com.br <sup>(2)</sup>Universidade Federal de Santa Maria (UFSM), Centro de Ciências Rurais, Departamento de Solos, Avenida Roraima, nº 1.000, Cidade Universitária, Camobi, CEP 97105-900 Santa Maria, RS, Brazil. E-mail: dalvan@ufsm.br, paulogubiani@gmail.com, rpmulazzani@gmail.com, somavillaa@gmail.com <sup>(3)</sup>UFSM, Departamento de Ciências Agronômicas e Ambientais, Campus Frederico Westphalen, Linha 7 de Setembro, BR 386, Km 40, CEP 98400-000 Frederico Westphalen, RS, Brazil. E-mail: vanderlei@ufsm.br

**Abstract** – The objective of this work was to evaluate the effect of different amounts of black oat (*Avena strigosa*) straw covering soil surface on soil temperature at different depths. The treatments consisted of 0, 3, 6, and 9 Mg ha<sup>-1</sup> straw. Soil temperature was measured hourly by a thermocouple inserted at different depths (0, 5, 15, 30, and 50 cm) and was used to adjust an equation correlating the temperature of covered soil with that of bare soil. With the correlations, it was possible to observe a point value of temperature (inversion temperature of straw effect), below which the presence of straw acts positively on the maintenance of soil temperature and above which the presence of straw acts negatively on soil heating.

**Index terms:** *Avena strigosa*, inversion temperature, mulch effect, no-tillage, plant residue, soil thermal regime.

### Alterações da temperatura em solo coberto de palha de aveia-preta

**Resumo** – O objetivo deste trabalho foi avaliar o efeito de diferentes quantidades de palha de aveia-preta (*Avena strigosa*) em cobertura do solo sobre a temperatura do solo em diferentes profundidades. Os tratamentos consistiram de 0, 3, 6 e 9 Mg ha<sup>-1</sup> de palha. A temperatura do solo foi medida a cada hora por meio de termopares inseridos em diferentes profundidades (0, 5, 15, 30 e 50 cm) e usada para ajustar uma equação que correlaciona a temperatura do solo coberto com a do solo descoberto. A partir dessas correlações, foi possível observar um valor pontual de temperatura (temperatura de inversão do efeito da palha), abaixo do qual a presença de palha atua positivamente na manutenção da temperatura do solo e acima do qual a presença de palha atua negativamente no aquecimento do solo.

**Termos para indexação:** *Avena strigosa*, temperatura de inversão, efeito da palha, plantio direto, resíduo vegetal, regime térmico do solo.

Worldwide, around 111 million hectares are cultivated under no-tillage system (Derpsch et al., 2010). In Brazil, 32 million hectares (FEBRAPDP, 2012) are planted under this direct-seeding system, for which soil cover by crop residues is recommended. The presence of straw cover affects soil temperature (Ts), which is attributed to the changes in the albedo surface and to the lower thermal conductivity of the straw layer compared with that of the soil.

Straw mulch has been shown to reduce Ts (Silva et al., 2006; Webler et al., 2016) and thermal amplitude (Dahiya et al., 2007; Furlani et al., 2008; Coelho et al., 2013). Ramakrishna et al. (2006) and Liu et al. (2014) found that, during cold periods, Ts of covered soil is

greater than that of bare soil; however, the inverse was observed during hot periods.

It should be noted that, in several studies, it may not have been possible to detect increases in Ts of mulching soil, because only daily or weekly Ts averages were measured. In fact, Ts of soil covered by straw may remain higher than that of bare soil at night or during periods of Ts depletion (Silva et al., 2006; Chen et al., 2007). Besides, most researches do not take into account the variation in straw amounts.

The objective of this work was to evaluate the effect of different amounts of black oat straw covering soil surface on soil temperature at different depths.

The study was carried out in the experimental area of Universidade Federal de Santa Maria, in the state of Rio Grande do Sul, Brazil (29°43'40"S, 53°43'11"W, at an average altitude of 101 m). The climate of the region is of the Cfa type according to Köppen (Peel et al., 2007). The soil was classified as an Argissolo Vermelho distrófico arênico, according to the Brazilian soil classification system (Santos et al., 2013), i.e., a Rhodic Paleudalf (Soil Survey Staff, 2014). The textural characterization of the soil, classified as sandy loam, showed: 665 g kg<sup>-1</sup> sand, 212 g kg<sup>-1</sup> silt, and 123 g kg<sup>-1</sup> clay in the 0–20-cm soil layer; 628 g kg<sup>-1</sup> sand, 244 g kg<sup>-1</sup> silt, and 128 g kg<sup>-1</sup> clay in the 20–37-cm layer; and 637 g kg<sup>-1</sup> sand, 251 g kg<sup>-1</sup> silt, and 112 g kg<sup>-1</sup> clay in the 37–50-cm layer.

The treatments consisted of different mulch amounts: 0 (bare soil), 3, 6, and 9 Mg ha<sup>-1</sup> dry mass of black oat (*Avena strigosa* Schreb.) straw. The straw was distributed on 6-m<sup>2</sup> (2x3 m) experimental units on 11/24/2014 and renewed (i.e., picked up and replaced by fresh dry straw) on January 12, March 11, April 24, June 13, August 13, and October 22, 2015. The straw was collected in an adjacent area where black oat was sown in May 2014; the whole plant was harvested in full bloom, dried under the sun, and stored in a covered place. The pre-emergent herbicide flumyazin was applied when the straw was renewed, in order to avoid the appearance of plants in the experimental units.

Ts was measured in the period from 12/1/2014 to 12/6/2015. In 2015, data recorded between October 7 and 22 were disregarded since intense winds moved the straw from soil surface. Ts was measured using the type T copper-constantan thermocouple. A polyethylene cap (3 cm of width and 0.6 mm of diameter), filled with epoxy resin to avoid metal alloy oxidation, was placed at the extremity of each sensor. The sensors were connected to a CR1000 datalogger (Campbell Scientific, Inc., Logan, UT, USA), programmed to perform instantaneous Ts measurements and to automatically record data every hour. The sensors were installed at a depth of 0 (soil surface), 5, 15, 30, and 50 cm in the central zone of the plot, in order to eliminate the effect of the adjacent treatments.

For each depth (0, 5, 15, 30, and 50 cm) and each treatment (0, 3, 6, and 9 Mg ha<sup>-1</sup>), Ts of soil covered by straw (Ts<sub>st</sub>) was correlated to the respective Ts of bare soil (Ts<sub>bs</sub>), both measured at exactly the same time. The linear regression equation,  $Ts_{st} = b_0 + b_1 Ts_{bs}$ , was adjusted

using the least squares method to estimate the  $b_0$  and  $b_1$  coefficients.

The  $b_0$  and  $b_1$  coefficients were compared with those of the equation of the 1:1 line, where  $b_1 = 1$  and  $b_0 = 0$ , indicating that  $Ts_{st} = Ts_{bs}$  and that there was no straw effect on Ts. The coefficients were compared by the t-test, considering the hypotheses  $b_0 \neq 0$  and  $b_1 \neq 1$ , at 5% probability of error. The intersection of the  $Ts_{st} = b_0 + b_1 Ts_{bs}$  line with the  $Ts_{st} = Ts_{bs}$  line, calculated by the equation  $Ts_{bs} = b_0 / (1 - b_1)$ , indicates the inversion set point of the mulch effect on Ts, i.e., when the temperature values for Ts of bare soil and of soil covered by straw were equal.

The  $b_1$  and  $b_0$  coefficients (Figure 1) were statistically different from 1 and 0, respectively, indicating that the presence of straw on soil surface modified Ts up to a depth of 50 cm. In all cases, the  $b_1$  coefficient remained lower than 1 and the  $b_0$  coefficient remained higher than 0. A  $b_0$  higher than 0 indicates a higher Ts in soil covered by straw at the beginning of the day, and a  $b_1$  lower than 1 indicates that the heating and cooling rates of soil with straw covering are lower than that of bare soil. In addition, the reflectivity and albedo of straw are greater than those of bare soil (Liu et al., 2014), which reduces the amount of energy available for heating the surface of mulched soil. In the straw mulch layer, a depletion of heat flux also occurs due to its low heat conductivity, which reduced both soil heating (during the day) and cooling (during the night) in depth. These results are in alignment with those described by Webler et al. (2016), indicating lower energy losses in covered soil.


The temperature at the intersection point of the Ts<sub>st</sub> regression line with the 1:1 line was referred to as the inversion temperature of straw effect (T<sub>inv</sub>). Therefore, when Ts<sub>bs</sub> is higher than T<sub>inv</sub>, covered soils have lower temperatures than bare soils; however, when Ts<sub>bs</sub> is lower than T<sub>inv</sub>, the opposite is observed (Figure 1). At the same depth, T<sub>inv</sub> showed variations of up to 0.5°C between treatments (Figure 1). On average, on soil surface, T<sub>inv</sub> was 19.6°C and, at 50 cm-depth, 16°C. Higher T<sub>inv</sub> values near soil surface may be associated with the larger temperature amplitudes of the surface layers. Studies on the effect of straw cover on Ts at a 5-cm depth showed Ts<sub>st</sub> lower than Ts<sub>bs</sub> when Ts<sub>bs</sub> was greater than 23°C (Furlani et al., 2008) and 20°C (Ribas et al., 2015). However, Almeida (2011) found Ts<sub>bs</sub> values lower than Ts<sub>st</sub> when Ts<sub>bs</sub> was 15.6°C.

The increment in the straw amount caused an increase in  $b_0$  and a decrease in  $b_1$  (Figure 1), indicating that the difference between  $T_{S_{bs}}$  and  $T_{S_{st}}$  is positively related to the amount of straw. The increase in straw amount reduced soil energy losses, resulting in a higher temperature for  $T_{S_{bs}}$  and a lower one for  $T_{inv}$ . However, if  $T_{S_{bs}}$  is greater than  $T_{inv}$ , the effect is the opposite.

The increase in straw amount reduced both the downward and upward heat flux densities by increasing the path of heat through the straw ( $\Delta Z$ ) as the straw layer became thicker. As a result, the heat propagation from straw surface to soil surface, and vice-versa, takes place with a small thermal gradient ( $\Delta T/\Delta Z$ ). Considering that the thermal conductivity ( $k$ ) of the medium (straw and air) is similar in all straw amounts, the Fourier equation,  $q = k(\Delta T/\Delta Z)$  (Dong et al., 2015), indicates a decrease in the heat flux density ( $q$ ) with the decrease in the  $\Delta T/\Delta Z$  thermal gradient.

The greater the depth, the lower the straw effect on  $T_s$  variation, i.e., the  $b_1$  angular coefficient is close to 1 and the  $b_0$  intersection value to 0. Consequently, the line that correlates  $T_{S_{st}}$  and  $T_{S_{bs}}$  is close to the 1:1 line, clearly indicating that changes in  $T_s$  caused by mulch decrease with depth. As a result, the straw cover effect on  $T_s$  was canceled out at depths below 50 cm.

According to several authors, mulch effects on soil temperatures vary with the presence of crop. Dalmago et al. (2004) and Silva et al. (2006) reported no effect of straw cover on soil temperature at 10 cm-depth under corn (*Zea mays* L.) and bean (*Phaseolus vulgaris* L.) vegetative canopies. In turn, Furlani et al. (2008), at 5-cm depth, found that soil temperature was not affected under the vegetative canopies of black oat and radish (*Raphanus sativus* L.). This indicates that the effect of mulch on temperatures at different soil depths depends on the conditions of the crop and on the interception of solar radiation. Since, in the present study, no plant was cultivated, solar radiation directly



**Figure 1.** Relationship between the temperatures of soil ( $T_{S_{st}}$ ) covered by 3 (I), 6 (II), and 9 (III) Mg ha<sup>-1</sup> straw and of bare soil ( $T_{S_{bs}}$ ) at the following soil depths: 0 cm (A), 5 cm (B), 15 cm (C), 30 cm (D), and 50 cm (E).  $T_{inv}$ , inversion temperature of straw effect.

impacted bare soil and the surface mulch layer. This might have resulted in higher thermal gradients in bare soil, compared with those in covered ones, consequently causing higher temperature variations in depth between treatments.

The obtained data showed that  $T_s$  was higher at the beginning of the day in soil covered by straw, but, that, throughout the day, heating penetration was dampened, compared with bare soil. The straw heats and cools the soil before and after bare soil reaches a given temperature, defined as the inversion temperature of straw effect. The inversion temperature decreases with depth; however, at the same depth, it is not affected by the amount of straw on soil surface. Black oat straw affects soil temperature up 50-cm depth, and the greater the amount of mulch, the more intense is the effect.

## References

- ALMEIDA, R.E. de. **Evaporação e temperatura em solos mantidos com diferentes quantidades de resíduos em superfície**. 2011. 116p. Tese (Doutorado) – Universidade Federal de Santa Maria, Santa Maria.
- CHEN, S.Y.; ZHANG, X.Y.; PEI, D.; SUN, H.Y.; CHEN, S.L. Effects of straw mulching on soil temperature, evaporation and yield of winter wheat: field experiments on the North China Plain. **Annals of Applied Biology**, v.150, p.261-268, 2007. DOI: 10.1111/j.1744-7348.2007.00144.x.
- COELHO, M.E.H.; FREITAS, F.C.L.; CUNHA, J.L.X.L.; SILVA, K.S.; GRANGEIRO, L.C.; OLIVEIRA, J.B. Coberturas do solo sobre a amplitude térmica e a produtividade de pimentão. **Planta Daninha**, v.31, p.369-378, 2013. DOI: 10.1590/S0100-83582013000200014.
- DAHIYA, R.; INGWERSEN, J.; STRECK, T. The effect of mulching and tillage on the water and temperature regimes of a loess soil: experimental findings and modeling. **Soil and Tillage Research**, v.96, p.52-63, 2007. DOI: 10.1016/j.still.2007.02.004.
- DALMAGO, G.A.; BERGAMASCHI, H.; COMIRAN, F.; BIANCHI, C.A.M.; BERGONCI J.I.; HECKLER, B.M.M. Soil temperature in maize crops as function of soil tillage systems. In: INTERNATIONAL SOIL CONSERVATION ORGANIZATION CONFERENCE, 13., 2004, Brisbane. **Conserving soil and water for society: sharing solutions: proceedings**. Brisbane: ISCO, 2004. 4p. ISCO 2004. Resumos expandidos.
- DERPSCH, R.; FRIEDRICH, T.; KASSAM, A.; HONGWEN, L. Current status of adoption of no-till farming in the world and some of its main benefits. **International Journal of Agricultural and Biological Engineering**, v.3, p.1-25, 2010. DOI: 10.3965/j.issn.1934-6344.2010.01.0-0.
- DONG, Y.; MCCARTNEY, J.S.; LU, N. Critical review of thermal conductivity models for unsaturated soils. **Geotechnical and Geological Engineering**, v.33, p. 207-221, 2015. DOI: 10.1007/s10706-015-9843-2.
- FEBRAPDP. Federação Brasileira de Plantio Direto na Palha. **Evolução da área cultivada no sistema plantio direto na palha – Brasil**. 2012. Available at: <[http://febrapdp.org.br/download/PD\\_Brasil\\_2013.I.pdf](http://febrapdp.org.br/download/PD_Brasil_2013.I.pdf)>. Accessed on: Dec. 23 2016.
- FURLANI, C.E.A.; GAMERO, C.A.; LEVIEN, R.; SILVA, R.P. da; CORTEZ, J.W. Temperatura do solo em função do preparo do solo e do manejo da cobertura de inverno. **Revista Brasileira de Ciência do Solo**, v.32, p.375-380, 2008. DOI: 10.1590/S0100-06832008000100035.
- LIU, Y.; WANG, J.; LIU, D.; LI, Z.; ZHANG, G.; TAO, Y.; XIE, J.; PAN, J.; CHEN, F. Straw mulching reduces the harmful effects of extreme hydrological and temperature conditions in citrus orchards. **PLoS ONE**, v.9, e87094, 2014. DOI: 10.1371/journal.pone.0087094.
- PEEL, M.C.; FINLAYSON, B.L.; MCMAHON, T.A. Updated world map of the Köppen-Geiger climate classification. **Hydrology and Earth System Science**, v.11, p.1633-1644, 2007. DOI: 10.5194/hess-11-1633-2007.
- RAMAKRISHNA, A.; TAM, H.M.; WANI, S.P.; LONG, T.D. Effect of mulch on soil temperature, moisture, weed infestation and yield of groundnut in northern Vietnam. **Field Crops Research**, v.95, p.115-125, 2006. DOI: 10.1016/j.fcr.2005.01.030.
- RIBAS, G.G.; STRECK, N.A.; SILVA, S.D. da; ROCHA, T.S.M. da; LANGNER, J.A. Temperatura do solo afetada pela irrigação e por diferentes coberturas. **Engenharia Agrícola**, v. 35, p.817-828, 2015. DOI: 10.1590/1809-4430-Eng.Agric.v35n5p817-828/2015.
- SANTOS, H.G. dos; JACOMINE, P.K.T.; ANJOS, L.H.C. dos; OLIVEIRA, V.A. de; LUMBRERAS, J.F.; COELHO, M.R.; ALMEIDA, J.A. de; CUNHA, T.J.F.; OLIVEIRA, J.B. de. **Sistema brasileiro de classificação de solos**. 3.ed. rev. e ampl. Brasília: Embrapa, 2013. 353p.
- SILVA, V.R. da; REICHERT, J.M.; REINERT, D.J. Variação na temperatura do solo em três sistemas de manejo na cultura do feijão. **Revista Brasileira de Ciência do Solo**, v.30, p.391-399, 2006. DOI: 10.1590/S0100-06832006000300001.
- SOIL SURVEY STAFF. **Keys to soil taxonomy**. 12<sup>th</sup> ed. Washington: USDA, 2014. 372p.
- WEBLER, G.; ROBERTI, D.R.; DIAZ, M.B.; TEISCHRIEB, C.C.; ZWIRTES, A.L.; REINERT, D.J. Efeitos de uma camada de palha no comportamento térmico do solo. **Ciência e Natura**, v.38, p.7-10, 2016. Edição especial. DOI: 10.5902/2179460X20381.

Received on January 7, 2017 and accepted on April 11, 2017