

# Study of metalloproteinases in the blood of goats experimentally infected with caprine encephalitis arthritis virus<sup>1</sup>

## Estudo de metaloproteínas no sangue de caprinos experimentalmente infectados pelo vírus da artrite encefalite caprina

Ylana Santos de Galiza<sup>2</sup>; Angela Maria Xavier Eloy<sup>3</sup>; Raymundo Rizaldo Pinheiro<sup>3</sup>; Renato Mesquita Peixoto<sup>4</sup>; Ana Milena César Lima<sup>5\*</sup>; Maria Luane da Silva Barroso<sup>6</sup>; Luzianna Macedo Fonseca<sup>7</sup>

### Highlights:

MMPs is a new approach to complement the diagnosis of caprine arthritis encephalitis.

Presence of proMMP-13 may be indicative of seroconversion in acute infection.

ProMMP-9 at post-seroconversion stage can facilitate control of the disease.

MMP-2 at post-seroconversion stage indicates the presence of infection.

### Abstract

Caprine arthritis encephalitis is a lentiviral disease that leads to considerable losses in goat farming. In the acute phase of viral infection, though antiviral antibodies are produced by the host's immune system, they are not sufficient to be detected by serological tests. Acute infections begin with an incubation period, during which the viral genome replicates and host innate responses are initiated. Matrix metalloproteinases (MMPs) are enzymes that play an important role in the physiological and pathological processes of tissue remodeling. The present study aimed to evaluate the expression of MMPs and their activity in the blood serum of male goats experimentally infected with caprine arthritis encephalitis virus (CAEV). Five dairy male goats, aged 3-4 years, were intravenously inoculated with CAEV Cork strain (titer:  $10^{5.6}$  TCID<sub>50</sub>/mL) after being tested negative for CAEV thrice at consecutive intervals of 30 days using western blot analysis and nested-PCR. The study included three stages: S1 or pre-infection stage; S2 or seroconversion stage, corresponding to the occurrence of first seroconversion; and S3 or post-seroconversion stage, corresponding to 23 weeks after seroconversion. Zymography was performed for the samples using gelatin zymography gels (12.5%), which were subjected to electrophoresis at 170V, 1A, and 300W for 50-70 min. The density of MMP-2 was found to be lower at

<sup>1</sup> Parte da Dissertação de Mestrado do primeiro autor.

<sup>2</sup> Discente do Curso de Doutorado do Programa de Pós-Graduação em Zootecnia, Universidade Federal do Ceará, UFC, Fortaleza, CE, Brasil. E-mail: ylana6@gmail.com

<sup>3</sup> Pesquisadores, Departamento de Sanidade Animal, Empresa Brasileira de Pesquisa Agropecuária, EMBRAPA CAPRINOS E OVINOS, Sobral, CE, Brasil. E-mail: angela.elay@embrapa.br; rizaldo.pinheiro@embrapa.br

<sup>4</sup> Dr., EMBRAPA CAPRINOS E OVINOS, Sobral, CE, Brasil. Bolsista de Desenvolvimento Científico Regional do Conselho Nacional de Desenvolvimento Científico e Tecnológico (DCR-CNPq/FUNCAP), nível C, Brasília, DF, Brasil. E-mail: renatomiraima@gmail.com

<sup>5</sup> Discente do Curso de Doutorado do Programa de Pós-Graduação em Ciência Animal, Universidade Federal do Piauí, UFPI, Teresina, PI, Brasil. E-mail: anamilenalima@yahoo.com.br

<sup>6</sup> M.e em Zootecnia, Universidade Estadual Vale do Acaraú, UVA, Sobral, CE, Brasil. E-mail: luanesilva62@hotmail.com

<sup>7</sup> Discente do Curso de Graduação em Medicina Veterinária, Centro Universitário INTA, UNINTA, Sobral, CE, Brasil. E-mail: luzianna.medicinavet@gmail.com

\* Author for correspondence

S1 (1456.20 pixels) than that at S2 and S3 (1943.80 and 2104.40 pixels, respectively) ( $P < 0.05$ ); and the density of MMP-9 was found to be lower at S3 (133.60 pixels) than that at S1 and S2 (359.60 and 370.60 pixels, respectively). The density of proMMP-2 was low at S1 and S3 (130.45 and 145.20 pixels, respectively). On the other hand, the density of proMMP-9 was statistically different between S1 and S3 (89.22 vs. 415.60 pixels). Both proMMP-2 and proMMP-9 were absent at S2. Thus, MMP-2 and MMP-9 exhibited opposite behaviors depending on the stage of infection. As the greatest activity of MMP-2 was detected at stage S3, we suggest that MMP-2 can be used as a biomarker for complementary diagnosis of acute CAEV infection. In addition, the presence of proMMP-13 can be used to indicate active viral infection.

**Key words:** Lentivirus. MMPs. Proteases. Zymography.

## Resumo

A encefalite por artrite caprina é uma doença lentiviral que leva a perdas consideráveis na criação de caprinos. Na fase aguda da infecção viral, embora os anticorpos antivirais sejam produzidos pelo sistema imunológico do hospedeiro, eles não são suficientes para serem detectados por testes sorológicos. As infecções agudas começam com um período de incubação, durante o qual o genoma viral se replica e as respostas inatas do hospedeiro são iniciadas. As metaloproteinases da matriz (MMPs) são enzimas que desempenham um papel importante nos processos fisiológicos e patológicos da remodelação tecidual. O presente estudo teve como objetivo avaliar a expressão de MMPs e sua atividade no soro sanguíneo de reprodutores caprinos infectados experimentalmente pelo vírus da encefalite por artrite caprina (CAEV). Cinco machos caprinos, com idades entre 3-4 anos, foram inoculadas por via intravenosa com a cepa de CAEV Cork (título:  $10^{5-6}$  TCID<sub>50</sub>/mL) após serem testados negativamente para CAEV três vezes em intervalos consecutivos de 30 dias usando a análise de western blot e nested-PCR. O estudo incluiu três etapas: estágio S1 ou pré-infecção; S2 ou estágio de soroconversão, correspondente à ocorrência de primeira soroconversão; e S3 ou pós-soroconversão, correspondendo a 23 semanas após a soroconversão. A zimografia foi realizada para as amostras utilizando gel de gelatina (12,5%), que foram submetidos à eletroforese em 170V, 1A e 300W por 50-70 min. Verificou-se que a densidade de MMP-2 era menor em S1 (1456,20 pixels) do que em S2 e S3 (1943,80 e 2104,40 pixels, respectivamente) ( $P < 0,05$ ); e a densidade de MMP-9 foi menor em S3 (133,60 pixels) do que em S1 e S2 (359,60 e 370,60 pixels, respectivamente). A densidade do proMMP-2 era baixa em S1 e S3 (130,45 e 145,20 pixels, respectivamente). Por outro lado, a densidade do proMMP-9 foi estatisticamente diferente entre S1 e S3 (89,22 vs. 415,60 pixels). ProMMP-2 e proMMP-9 estavam ausentes no S2. Assim, MMP-2 e MMP-9 exibiram comportamentos opostos, dependendo do estágio da infecção. Como a maior atividade da MMP-2 foi detectada no estágio S3, sugerimos que a MMP-2 possa ser usada como biomarcador para o diagnóstico complementar de infecção aguda por CAEV. Além disso, a presença de proMMP-13 pode ser usada para indicar infecção viral ativa.

**Palavras-chave:** Lentivírus. MMPs. Proteases. Zimografia.

## Introduction

Caprine arthritis encephalitis leads to great economic losses in goat production, particularly when breeders and matrices of high genetic value are affected by the disease; importantly, asymptomatic animals represent a serious threat to herd health (Pinheiro, Gouveia, Alves, & Andrioli, 2004). Although animals infected with caprine arthritis encephalitis virus (CAEV) develop specific

cellular and humoral immune responses against the virus, they are not capable of preventing viral transmission among animals (Dawson, 1987). Some infected animals exhibit delayed the production of detectable antibodies (Rimstad et al., 1993), which leads to false negative results in serological tests. The false negative results can also be attributed to the antigenic variation between the viral strain used in the test and the circulating strain (Jones, 2014). In addition, production of antibodies and their

concentration in the blood are influenced by several factors, including stress, nutritional condition, age of the animal, and presence of other diseases (Franke, 1998).

The time interval between infection and seroconversion varies among animals, ranging from weeks to months or even years, with the possibility of seroreversion making it even more difficult to control the disease (Hanson, Hydbring, & Olsson, 1996). In most cases, at third week post infection, immune response against the capsid protein (p25 or p28) (Houwens & Nauta, 1989) can be detected; and at fifth week post infection, antibodies against other proteins can be detected (Concha-Bermejillo, Brodie, Magnus-Corral, Brown, & DeMartini, 1995).

Proteomics has emerged as an advanced tool to circumvent and promote effective control of infections because it facilitates the study of molecules involved in innate immune response to bacterial and viral infections (Aslam, Basit, Nisar, Khurshid, & Rasool, 2017).

Among the expressed proteases, matrix metalloproteinases (MMPs), which play important roles as inflammatory mediators, enable the monitoring and prognosis of diseases and can serve as potential biomarkers for diagnosis of viral infections, subsequently facilitating the discovery of antiviral compounds (Pereira, 2014). MMPs are calcium-dependent zinc-containing endopeptidases secreted by polymorphonuclear leukocytes, keratinocytes, monocytes, macrophages, fibroblasts, and mesenchymal cells as proenzymes (Navarro, Nelson, Silva, & Freitas, 2006). A total of 24 MMPs have been identified in mammals, and they are classified into five subfamilies: collagenases (MMP-1, MMP-8, and MMP-13), gelatinases (MMP-2 and MMP-9), stromelysins (MMP-3 and MMP-10), elastases (MMP-7 and MMP-12), and membrane-like MMPs (MT-MMP-1 to MT-MMP-5) (Visse

& Nagase, 2003; Lopez-Avila & Spencer, 2008). MMPs participate in several biological activities, including determination of the structure of extracellular matrix, embryonic development, implantation of blastocysts, organ morphogenesis, development of the nervous system, ovulation, cervical dilation, postpartum uterine regression, development and remodeling of oral tissue, tissue healing, angiogenesis, and apoptosis (Verma & Hansch, 2007). Therefore, MMPs are recognized as important regulatory enzymes exhibiting beneficial and harmful activities in the inflammatory pathways, where they may contribute to the regulation of host defense and inflammatory diseases (Concha, 2010). To evaluate the active and latent forms of proteolytic enzymes and to determine the number and approximate molecular mass of enzymes in biological samples, zymography, a method based on the degradation of a copolymerized substrate, is commonly employed (Eloy, Bezerra, Pinheiro, & Andrioli, 2015).

In medicine, studies related to the application of MMPs in monitoring, control, and diagnosis of bacterial and viral infections, diabetes, cancer and its progression, and other diseases, are at an advanced stage. On the contrary, in veterinary medicine, studies on MMPs are still incipient, especially in goats. Studies related to the role of MMPs in goat diseases are scarce. However, Bezerra et al. (2015) described the high activity of MMPs (proMMP-2 and MMP-2) in goats naturally infected with CAEV, both in the presence and absence of joint symptoms. Considering the important roles of MMPs in the physiological processes of animals, particularly CAEV-infected animals, and the lack of information regarding the performance of these enzymes in goats experimentally infected with CAEV, the present study aimed to acquire a comprehensive understanding of the functions of MMPs in the blood serum of male goats experimentally infected with CAEV.

## Materials and Methods

A total of five males goats Saanen and Anglo-Nubian goats, aged 3-4 years, belonging to the herd of EMBRAPA Goats and Sheep, located in the municipality of Sobral, Ceará, Brazil, were used in the present study. The animals were tested for CAEV thrice at consecutive intervals of 30 days using western blotting (WB) and nested-PCR, and it was observed that all animals tested negative for CAEV. All animals were healthy according to a general clinical examination performed following the recommendations of Diffay, Mckenzie, Wolf and Pugh (2005). The animals were fed elephant grass (*Pennisetum purpureum*) supplemented with 300 g concentrate containing 70 % corn, 27 % soybean meal, 2 % limestone, and 1 % mineral salt *ad libitum*. The study was carried out in accordance with the ethical principles of animal experimentation and according to the protocol submitted to and subsequently approved by the Institutional Animal Care and Use Committee of the Universidade Estadual Vale do Acaraú (UVA) on December 17, 2014, under protocol number 001.04.013.UVA.505.01.

The five CAEV-free animals were inoculated with 1 mL of minimum essential medium containing CAEV Cork strain (titer:  $10^{5-6}$  TCID<sub>50</sub>/mL) intravenously, followed by weekly follow-up with WB until seroconversion and monthly follow-up thereafter. Thus, the study included three stages: S1 or pre-infection stage, corresponding to one week before viral inoculation; S2 or seroconversion stage, corresponding to third week after viral inoculation i.e., the occurrence of first seroconversion; and S3 or post-seroconversion stage, corresponding to 23 weeks after seroconversion.

Blood samples were collected by jugular venipuncture using a vacutainer® system in 10 mL tubes without anticoagulant, and centrifuged at 3000 g for 10 min for separation of serum. The serum was

stored in a 2.5 mL microtube at 20 °C until further analysis.

Protein concentration was determined using Bradford method (Bradford, 1976). Zymography was performed using 12.5 % polyacrylamide gels copolymerized with sodium dodecyl sulfate (SDS), with gelatin (2 mg/mL gelatin) as the substrate, according to the protocol of Kupai et al. (2010). Electrophoresis was carried out at 170 V, 1 A, and 300 W for 50-70 min (Laemmli, 1970). The gels were prepared in duplicates for each stage (S1, S2, and S3), and were scanned and analyzed using Doc-IT® LS Image Analysis Software 6.0. The LMW-SDS marker kit (GE Healthcare, Chicago, IL, United States), containing phosphorylase B (97.4 kDa), albumin (67 kDa), ovalbumin (43 kDa), carbonic anhydrase (30 kDa), soybean trypsin inhibitor (20.1 kDa), and lactalbumin (14.4 kDa), was used as protein molecular weight standard. Active MMP-9 (86 kDa) and MMP-2 (66 kDa) and their respective latent forms, proMMP-9 (92 kDa) and pro proMMP-2 (72 kDa), as well as other MMPs are expected to have an affinity towards gelatin. Recently, these MMPS were found in the blood serum and seminal plasma samples of goats chronically infected with CAEV. (Bezerra et al., 2015, 2017; Pinto et al., 2019; Santos et al., 2019).

Statistical analysis for this study was descriptive, analyzing the presence or absence of bands corresponding to the respective enzymes in the gels. The mean volume and intensity of each enzyme was calculated using Gel Analyzer version 2010®. The band intensity data were subjected to Shapiro-Wilk and Bartlett tests for verification of the assumptions of normality and homogeneity of variances. Logarithmic transformation of base 10 was applied to guarantee such assumptions. F test was performed using analysis of variance, followed by Tukey's posthoc test, using a 5 % significance level. All statistical analyses were performed using the R software.

## Results and Discussion

Seroconversion was detected using WB in all animals at third week post infection. In the post-seroconversion stage i.e., 23 weeks after infection, only one animal tested negative for CAEV, while the others remained seropositive (Table 1). Lara, Birgel, Fernandes and Birgel (2003) injected infected goat kids with blood and colostrum from infected animals, and found that the titer of specific antibodies reached a maximum level between 45 and 60 days post-infection, and decreased thereafter.

Our results corroborate the work of Spivak et al. (2010), who demonstrated that seroconversion occurred in human immunodeficiency virus-infected individuals within 3 weeks to one year. The variation in the seroconversion period can be attributed to the tropism of CAEV to monocytes and macrophages, which play important roles in viral replication and diffusion (Blacklaws, 2012). According to Pugh (2004), restricted replication allows the virus to remain latent in host monocytes, making them go undetected by the immune system.


**Table 1**  
Results of western blot analysis for detection of CAEV at pre-infection (one week before infection), seroconversion (third week post infection), and post-seroconversion (twenty-third week post infection) stages in the blood serum of goats experimentally infected with CAEV

Stage	Animals				
	1	2	3	4	5
Pre-infection (S1)	-	-	-	-	-
Seroconversion (S2)	+	+	+	+	+
Post-seroconversion (S3)	+	-	+	+	+

(+) seropositive for CAEV; (-) seronegative for CAEV.

At S1, latent proMMP-2 (72-75 kDa) and proMMP-9 (92 kDa) as well as active MMP-2 (64-66 kDa) and MMP-9 (80-84 kDa) were detected; at S2, only active MMP-2 and MMP-9 and latent proMMP-13 (48 kDa) were detected; and at S3,

the MMP profile was similar to that observed at S1 (Figure 1). High molecular weight proteins were detected in all animals; however, these were not the focus of this study.



**Figure 1.** Enzymatic activity of metalloproteinases in the blood serum of goats experimentally infected with caprine arthritis encephalitis virus. Animals (1-5); (A) pre-infection, (B) seroconversion, and (C) post-seroconversion.

Bezerra et al. (2015) found that the activity of proMMP-2 ( $P = 0.0025$ ) and MMP-2 ( $P = 0.0001$ ) was greater in breeders naturally infected with CAEV than that in animals of the control group (negative), suggesting that proteases play an important role in chronic inflammatory processes in goats. An increase in the activity of these enzymes and a greater number of peaks have also been demonstrated among seropositive animals. Using a two-dimensional electrophoresis technique and matrix assisted laser desorption/ionization-time of flight (MALDI-TOF) mass spectrometry, Bezerra et al. (2017) identified arisultadase A, a protease involved in the regulation of sulfate metabolism and subsequent control of viral infections, in the seminal plasma of goat breeders infected with CAEV.

MMPs are secreted as proenzymes or zymogens, therefore, they must be proteolytically activated by removal of the propeptide. This process is accompanied by breaking of the bond between the cysteine thiol group in the propeptide and the catalytic zinc in the peptide (Geurts, Opdenakker, & Steen, 2012; Mannello & Medda, 2012).

Analysis of band intensity in pixels (Figure 1) revealed that the density of MMP2 was greater at S3 than that at S1 and S2, suggesting a positive regulation. On the contrary, the density of MMP-9 was lesser at S1 and S3 than that at S2, suggesting a negative regulation.

Latent proMMP-2 and proMMP-9 were produced only at S1 and S3. Considering S1, this is possibly because they are required for the physiological

processes of healthy animals but at low activities (130.45 vs. 89.22 pixels). At S2, the latent forms were transformed into their active forms i.e., MMP-2 and MMP-9; and at this stage, MMP-2 had greater activity (1943.80 pixels) than MMP-9 (370.60 pixels). The presence of proMMP-13 in all animals undergoing seroconversion indicates active viral infection.

Thus, MMP-2 and MMP-9 exhibited opposite behaviors according to the stage of infection. In addition, MMP-2 exhibited a difference ( $P < 0.05$ ) between S1 and S3, while MMP-9 exhibited no statistical difference between the two stages. In healthy animals, the density of proMMP-9 was below 90 pixels at S1 and above 400 pixels at S3 ( $P < 0.05$ ). ProMMP-2 did not present statistically significant differences between S1 (130 pixels) and S3 (145.20 pixels) (Table 2).

**Table 2**

**MMP density (mean  $\pm$  SD) observed at pre-infection (one week before infection), seroconversion (third week post infection), and post-seroconversion (twenty-third week post infection) stages in the blood serum samples of goats experimentally infected with CAEV using zymography**

Treatment	MMP density (Pixels)			
	MMP-2	MMP-9	ProMMP-2	ProMMP-9
Pre-infection (S1)	1456.20 <sup>b</sup> $\pm$ 236.8	359.60 <sup>a</sup> $\pm$ 209.1	130.45 <sup>a</sup> $\pm$ 73	89.22 <sup>b</sup> $\pm$ 51.6
Seroconversion (S2)	1943.80 <sup>a</sup> $\pm$ 242	370.60 <sup>a</sup> $\pm$ 247.8	Absent	Absent
Post-seroconversion (S3)	2104.40 <sup>a</sup> $\pm$ 139.5	133.00 <sup>a</sup> $\pm$ 50.7	145.20 <sup>a</sup> $\pm$ 42.3	415.60 <sup>a</sup> $\pm$ 114.7

Different lower-case superscript letters in the same column indicate that the values differ statistically at 5 % significance.

The low intensity of most MMPs at S1 indicates that they are present in healthy animals and perform physiological functions inherent to the organism's development. Proteolytic enzymes are regulated by different mechanisms and are responsible for the degradation and remodeling of extracellular matrix components, which are involved in normal physiological processes and are associated with various diseases (Matrisian, 1990; Woessner, 1991).

The proteases MMP-2, MMP-9, and proMMP-13 were found in all animals at S2. The absence of latent proMMP-2 and proMMP-9 at S2 suggests that they were transformed into their respective active forms i.e., MMP-2 and MMP-9 due to immune response induced by the virus.

Elkington, White, Addington-Hall, Higgs and Edmonds (2005) demonstrated that MMP-2 and MMP-9 are required for the migration of

inflammatory cells *in vitro*. ProMMP-13 is required for the activation of most of the MMPs.

It is worth mentioning that CAEV infection induces a strong humoral and cellular immune response (Lofstedt, 2014). Owing to the role of MMPs in inflammatory processes, it is suggested that enzyme activity of MMPs in the blood serum can serve as a marker of recent infection in animals. The presence of MMPs in the blood serum is attributed to the secretion of these enzymes by components of the circulatory and immune systems, including neutrophils, monocytes, macrophages, and fibroblasts, as well as by tumor cells in response to a variety of stimuli (Woessner, 1991). MMPs play a fundamental role in the inflammatory process and are involved in the pathophysiological remodeling of the vascular wall (Kuzuya & Iguchi, 2003; Hu, Steen, Sang, & Opendakker, 2007).

Studies have revealed that MMP-9 is responsible for the release of the biologically active form of vascular endothelial growth factor, which plays an essential role in angiogenesis (Bergers et al., 2000; Klein & Bischoff, 2011). Moreover, MMP-9 has the ability to proteolytically degrade the basement membrane, which indicates the important role of MMP-9 in the formation of new blood vessels (Klein & Bischoff, 2011). MMP-9 is also expressed in conditions that require tissue remodeling, such as wound healing and tumor invasion (Stamenkovic, 2000; Klein & Bischoff, 2011). Bezerra et al. (2015) reported that the levels of proMMP-9 and MMP-9 were higher, though variable, in the seropositive group than those in the control (uninfected) group. MMP-2 can degrade gelatin, collagens (type I, IV, and V), elastin, and vitronectin, and is widely expressed by cells of the connective tissue (Stamenkovic, 2000). In addition, MMP-2 is involved in cellular migration, and is responsible for direct degradation of the basement membrane (Klein & Bischoff, 2011). Though both these MMPs work in different ways, they complement each other.

The presence of proMMP-13 in all animals at S2 suggests that it can be used as a biomarker of active CAEV infection. ProMMP-13 is involved in the degradation of proteoglycans and collagens, which are the main components of cartilages (Takahashi, Sasaki, Tsouderos, & Suda, 2003). It has been reported that patients with joint cartilage destruction exhibit high MMP-13 expression (Roach et al., 2005), suggesting the role of MMP-13 in cartilage degradation (Wang et al., 2013). Knäuper, López-Otín, Smith, Knight, & Murphy et al. (1996, 1997) observed that MMP-13 is overexpressed in joints and articular cartilage in patients with early osteoarthritis but hardly detected in normal adult tissues. MMP-13 can modify a large amount of extracellular proteins and cause degradation and remodeling of damaged tissues, acting mainly on the layers of synovial lining and synovial tissues (Iwamoto et al., 2011). In addition, MMP-13 is involved in several diseases, including rheumatoid

arthritis (Ye et al., 2007), coronary artery disease (Vasku et al., 2012), and abdominal aortic disease (Saracini et al., 2012).

Knäuper et al. (2002) demonstrated that, in fibrosarcoma cell line (HT1080), proMMP-13 was activated to a 48-kDa active form i.e., MMP-13 via a 56-kDa intermediate. In the present study, the 48-kDa form of MMP-13 was detected at S2, suggesting that it is the active form in animals infected with CAEV. In addition, MMP-9 was expressed at a high level at S2, probably because MMP-13 induced greater activation of this protease. *In vitro* studies have shown that MMP-13 induces activation of proMMP-9 and self-activation of MMP-13. (Hernández Rios et al., 2009). These results suggest that active MMP-13 plays an important role in the activation of other MMPs, and is crucial for bone metabolism and homeostasis (Vincenti & Brinckerhoff, 2002).

At S3, MMP-2 and MMP-9, together with their respective latent forms, were detected. It has been suggested that the expression of these enzymes varies depending on the type and stage of disease, presence of neoplasms and inflammations, and cell type (Parks & Mecham, 1998). Thus, these proteases occur naturally in healthy animals as well as in infected animals; however, their expression is increased in infected animals to facilitate control of the disease. The absence of proMMP-13 at S3 is probably because after the first seroconversion, when there is a decrease in antibody production, activation of proteases by proMMP-13 is not required. The presence of proMMP-13 at S2 stage active viral infection. In addition, though the animals tested negative for CAEV at S3, the MMP profile at S3 was similar to that at S2.

The observed variations in the MMP profile can be attributed to the fact that enzyme profiles vary according to the type and stage of disease, presence of neoplasms and inflammation, and cell type; as the extracellular matrix, present in all tissues, requires MMPs for physiological processes and tissue remodeling (Yo & Werb, 1998).

## Conclusion

At all stages (S1, S2 and S3), MMP-2 exhibited greater activity than MMP-9. The greatest activity of MMP-2 was observed in S3, suggesting that MMP-2 can be used as in monitoring infection. In addition, proMMP-13, detected at seroconversion, indicates active viral infection. MMPs can diagnose viral infections even during viral latency, when serological tests show negative results. This is the first study reporting the expression profile of MMPs in male goats with experimentally infected with CAEV.

## Acknowledgments

The authors thank the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for funding this research, Fundação Cearense de Apoio ao Desenvolvimento Científico e Tecnológico (FUNCAP) and Embrapa Caprinos e Ovinos (CNPACO) for providing the physical space and materials necessary for conducting this study.

## References

- Aslam, B., Basit, M., Nisar, M. A., Khurshid, M., & Rasool, M. H. (2017). Proteomics: technologies and their applications. *Journal of Chromatographic Science*, 55(2), 182-196. doi: 10.1093/chromsci/bmw 167
- Bergers, G., Brekken, R., McMahon, G., Vu, T. H., Itoh, T., Tamaki, K.,... Hanahan, D. (2000). Matrix metalloproteinase-9 triggers the angiogenic switch during carcinogenesis. *Nature Cell Biology*, 2(10), 737-744. doi: 10.1038/35036374
- Bezerra, R. Q., Jr., Eloy, A. M. X., Furtado, J. R., Pinheiro, R. R., Andrioli, A. P., Moreno, F. B.,... Teixeira, M. F. S. (2017). A panel of protein candidates for comprehensive study of Caprine Arthritis Encephalitis (CAE) infection. *Tropical Animal Health and Production*, 50(1) 43-48, 2017. doi: 10.1007/s11250-017-1398-1
- Bezerra, R. Q., Jr., Eloy, A. M. X., Perreira, E. P., Furtado, J. R., Souza, K. C. S., Lima, A. R.,... Teixeira, M. F. S. (2015). Avaliação das metaloproteinases de matriz no sangue de reprodutores caprinos naturalmente infectados com artrite encefalite caprina na região Semiárida do Brasil. *Acta Scientiae Veterinariae*, 43(1258), 1-7. Recuperado de <http://www.ufrgs.br/actavet/43/PUB%201258.pdf>
- Blacklaws, B. A. (2012). Small ruminant lentiviruses: immunopathogenesis of visna-maedi and caprine arthritis and encephalitis virus. *Comparative Immunology, Microbiology and Infectious Diseases*, 35(3), 259-269. doi: 10.1016/j.cimid.2011.12.003
- Bradford, M. M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*, 72(1-2), 248-254. doi: 10.1016/0003-2697(76)90527-3
- Concha, H. A. R. (2010). *Papel das metaloproteinases de matriz nas alterações da barreira hematoencefálica em ratos submetidos à sepse severa*. Dissertação de mestrado, Ciências da Saúde, Universidade do Extremo Sul Catarinense, Criciúma, RS, Brasil. Recuperado de <http://www.bib.unesc.net/pergamum/biblioteca/index.php?codAcervo=99541>
- Concha-Bermejillo, A. de la., Brodie, S. J., Magnus-Corral, S., Browen, R. A. & DeMartini, J. C. (1995). Pathologic and serologic response to isogenic twin lambs to phenotypically distinct lentiviruses. *Journal of Acquired Immune Deficiency Syndromes and Retrovirology*, 8(2), 116-123. doi: 10.1097/00042560-199502000-00002
- Dawson, M. (1987). Pathogenesis of maedi-visna. *Veterinary Record*, 120(19), 451-454. doi: 10.1136/vr.120.19.451
- Diffay, B. C., McKenzie, D., Wolf, C., & Pugh, D. G. (2005). Abordagem e exame de ovinos e caprinos. In D. G. Pugh (Eds.), *Clínica de ovinos e caprinos* (pp. 1-19). São Paulo, SP: Roca.
- Elkington, H., White, P., Addington-Hall, J., Higgs, R., & Edmonds, P. (2005). The healthcare needs of chronic obstructive pulmonary disease patients in the last year of life. *Palliative Medicine*, 19(6), 485-491. doi: 10.1191/0269216305pm1056oa
- Eloy, A. M. X., Bezerra, R. Q., Jr., Pinheiro, R. R., & Andrioli, A. (2015). *Técnica Zimográfica como método para monitoramento da Artrite Encefalite Caprina (CAE)*. (Comunicado Técnico, 155). Sobral, CE: Embrapa Caprinos e Ovinos, CNPC. 1-6. Recuperado de <http://ainfo.cnptia.embrapa.br/digital/bitstream/item/146807/1/CNPC-2015-Cot155.pdf>
- Franke, C. R. (1998). *Controle sanitário da artrite-encefalite caprina (CAE)*. Salvador, BA: EDUFBA.

- Geurts, N., Opendakker, G., & Steen, P. E. van den. (2012). Matrix Metalloproteinases as therapeutic targets in protozoan parasitic infections. *Pharmacology & therapeutics*, 133(3), 257-279. doi: 10.1016/j.pharmthera.2011.11.008
- Hanson, J., Hydrbring, E., & Olsson, K. (1996). A long term study of goats naturally infected with caprine arthritis-encephalitis virus. *Acta Veterinaria Scandinavica*, 37(1), 31-39. Recuperado de <http://pubmed.ncbi.nlm.nih.gov/8659344/>
- Hernández Rios, M., Sorsa T., Obrego'n, F., Tervahartiala, T., Valenzuela, M. A., Pozo, P.,... Gamonal, J. (2009). Proteolytic roles of matrix metalloproteinases (MMP)-13 during progression of chronic periodontitis: initial evidence for MMP-13/MMP-9 activation cascade. *Journal of Clinical Periodontology*, 36(12), 1011-1017. doi: 10.1111/j.1600-051X.2009.01488.x
- Houwers, D. J., & Nauta, I. M. (1989). Immunoblot analysis of the antibody response to ovine lentivirus infections. *Veterinary Microbiology*, 19(2), 127-139. doi: 10.1016/0378-1135(89)90078-3
- Hu, J., Steen, P. E. van den., Sang, Q. X. A., & Opendakker, G. (2007). Matrix metalloproteinase inhibitors as therapy for inflammatory and vascular diseases. *Nature Reviews Drug Discovery*, 6(6), 480-498. doi: 10.1038/nrd2308
- Iwamoto, N., Kawakami, A., Arima, K., Tamai, M., Nakamura, H., Kawashiri, S.,... Eguchi, K. (2011). Contribution of an adenine to guanine single nucleotide polymorphism of the matrix metalloproteinase-13 (MMP-13) -77 promoter region to the production of anticyclic citrullinated peptide antibodies in patients with HLA-DRB1\*shared epitope-negative rheumatoid arthritis. *Journal of Modern Rheumatology*, 21(3), 240-243. doi: 10.1007/s10165-010-0375-6
- Jones, B. T. (2014). *The current prevalence of caprine arthritis-encephalitis virus in midwestern goat herds*. Doctoral thesis, University of Nebraska, Lincoln, NE, USA. Retrieved from <http://digitalcommons.unl.edu/cgi/viewcontent.cgi?article=1014&context=vetscidiss>
- Klein, T., & Bischoff, R. (2011). Physiology and pathophysiology of matrix metalloproteinases. *Amino Acids*, 41(2), 271-290. doi: 10.1007/s00726-010-0689-x
- Knäuper, V., Bailey, L., Worley, J. R., Soloway, P., Patterson, M. L., & Murphy, G. (2002). Cellular activation of pro-MMP-13 by MTI-MMP depends on the C-terminal domain of MMP-13. *FEBS Letters*, 532(1-2), 127-130. doi: 10.1016/s0014-5793(02)03654-2
- Knäuper, V., Cowell, S., Smith, B., Lopez-Otin, C., O'Shea, M., & Morris, H. (1997). The role of the C-terminal domain of human collagenase-3 (MMP-13) in the activation of pro collagenase-3, substrate specificity, and tissue inhibitor of metalloproteinase interaction. *Journal of Biological Chemistry*, 272(12), 7608-7616. doi: 10.1074/jbc.272.12.7608
- Knäuper, V., López-Otín, C., Smith, B., Knight, G., & Murphy, G. (1996). Biochemical characterization of human collagenase-3. *Journal of Biological Chemistry*, 271(3), 1544-1550. doi: 10.1074/jbc.271.3.1544
- Kupai, K., Szucs, G., Cseh, S., Hajdu, I., Csonka, C., Csont, T., & Ferdinandy, P. (2010). Matrix metalloproteinase activity assays: Importance of zymography. *Journal of Pharmacology and Toxicology Methods*, 61(2), 205-209. doi: 10.1016/j.vascn.2010.02.011
- Kuzuya, M., & Iguchi, A. (2003). Role of matrix metalloproteinases in vascular remodeling. *Journal of Atherosclerosis and Thrombosis*, 10(5), 275-282. doi: 10.5551/jat.10.275
- Laemmli, U. K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*, 227(5259), 680-685. doi: 10.1038/227680a0
- Lara, M. C. C. S. H., Birgel, E. H., Jr., Fernandes, M. A., & Birgel, E. H. (2003). Infecção experimental do vírus da Artrite Encefalite dos caprinos em cabritos. *Arquivos do Instituto Biológico*, 70(1), 51-54. Recuperado de [http://www.biologico.agricultura.sp.gov.br/uploads/docs/arq/V70\\_1/lara.pdf](http://www.biologico.agricultura.sp.gov.br/uploads/docs/arq/V70_1/lara.pdf)
- Lofstedt, J. (2014). *Overview of caprine arthritis and encephalitis*. Retrieved from <http://www.msddvetmanual.com/generalized-conditions/capriner-arthritis-and-encephalitis/overview-of-capriner-arthritis-and-encephalitis>
- Lopez-Avila, V., & Spencer, J. V. (2008). Methods for detection of matrix metalloproteinases as biomarkers in cardiovascular disease. *Clinical Medicine Insights: Cardiology*, 2(1), 75-87. doi: 10.4137/CMC.S484
- Mannello, F., & Medda, V. (2012). Nuclear localization of matrix metalloproteinases. *Progress in Histochemistry and Cytochemistry*, 47(1), 27-58. doi: 10.1016/j.proghi.2011.12.002
- Matrisian, L. M. (1990). Metalloproteinases and their inhibitors in matrix remodeling. *Trends in Genetics*, 6(4), 121-125. doi: 10.1016/0168-9525(90)90126-q

- Navarro, V. P., Nelson, P., F., Silva, L. A. B., & Freitas, A. C. A. (2006). Participação das metaloproteínas da matriz nos processos fisiopatológicos da cavidade bucal. *Revista de Odontologia da UNESP, Araraquara*, 35(4), 233-238. Recuperado de <http://www.revodontolunesp.com.br/article/588017de7f8c9d0a098b4951>
- Parks, W. C., & Mecham, R. P. (1998). *Matrix metalloproteinases*. San Diego: Academic Press.
- Pereira, P. E. (2014). *Caracterização das metaloproteínas (MMPs) no plasma seminal de caprinos sadios e infectados pelo vírus da artrite encefalite caprina (CAE)*. Dissertação de mestrado, Universidade Estadual Vale do Acaraú, Sobral, CE, Brasil. Recuperado de <http://www.alice.cnptia.embrapa.br/alice/bitstream/doc/1025854/1/CNPC2014Caracterizacaodasmetaloproteinades.pdf>
- Pinheiro, R. R., Gouveia, A. M. G., Alves, F. S. F., & Andrioli, A. (2004). Perfil de propriedades no estado do Ceará relacionado à presença do lentivírus caprino. *Ciência Animal*, 14(1), 29-37. Recuperado de <http://www.uece.br/cienciaanimal/dmdocuments/Artigo3.2004.1.pdf>
- Pinto, T. M. F., Moreira, R. F., Matos, M. N. C., Soares, V. V. M., Aguiar, M. V. A., Aragão, P. T. T. D. de, ... Cunha, R. M. S. (2019). Evaluation of the proteomic profiles of ejaculated spermatozoa from Saanen bucks (*Capra hircus*). *Animal Reproduction*, 16(4), 902-913. doi: 10.21451/1984-3143-ar2019-0001
- Pugh, D. C. (2004). *Clínica de ovinos e caprinos*. São Paulo, SP: Roca.
- Rimstad, E., East, N. E., Torten, M., Higgins, J., DeRock, E., & Pedersen, N. C. (1993). Delayed seroconversion following naturally acquired caprine arthritis-encephalitis virus infection in goats. *American Journal of Veterinary Research*, 54(11), 1858-1862. Retrieved from <http://pubmed.ncbi.nlm.nih.gov/8291763/>
- Roach, H. I., Yamada, N., Cheung, K. S., Tilley, S., Clarke, N. M., Oreffo, R. O., ... Bronner, F. (2005). Association between the abnormal expression of matrix-degrading enzymes by human osteoarthritic chondrocytes and demethylation of specific CpG sites in the promoter regions. *Arthritis & Rheumatology*, 52(10), 3110-3124. doi: 3110-3124. 10.1002/art.21300
- Santos, K. F. dos, Eloy, A. M. X., Matos, M. N. C., Peixoto, R. M., Aragão, P. T. T. D., Pinheiro, R. R., & Cunha, R. M. S. da. (2019). Use of proteomics in the study of the acute phase of caprine arthritis encephalitis in seminal plasma. *Small Ruminant Research*, 181, 39-44. doi: 10.1016/j.smallrumres.2019.10.004
- Saracini, C., Bolli, P., Sticchi, E., Pratesi, G., Pulli, R., Sofi, F., ... Giusti, B. (2012). Polymorphisms of genes involved in extracellular matrix remodeling and abdominal aortic aneurysm. *Journal of Vascular Surgery*, 55(1), 171-179. doi: 10.1016/j.jvs.2011.07.051
- Spivak, A. M., Brennan, T. P., O'Connell, K. A., Sydnor, E., Williams, T. M., Siliciano, R. F., ... Blankson, J. N. (2010). A case of seronegative HIV-1 infection. *The Journal of Infectious Diseases*, 201(3), 341-345. doi: 10.1086/649822
- Stamenkovic, I. (2000). Matriz metaloproteínas in tumor invasion and metastasis. *Seminars in Cancer Biology*, 10(6), 415-433. doi: 10.1006/scbi.2000.0379
- Takahashi, N., Sasaki, T., Tsouderos, Y., & Suda, T. (2003). S 12911-2 inhibits osteoclastic bone resorption in vitro. *Journal of Bone Mineral Research*, 18(6), 1082-1087. doi: 10.1359/jbmr.2003.18.6.1082
- Vasku, A., Meluzin, J., Blahák, J., Kincl, V., Goldbergová, M. P., Sitar, J., ... Vítovec, J. (2012). Matrix metalloproteinase 13 genotype in rs640198 polymorphism is associated with severe coronary artery disease. *Diseases Markers*, 33(1), 43-49. doi: 10.3233 / DMA-2012-0902
- Verma, R. P., & Hansch, C. (2007). Matrix metalloproteinases (MMPs): chemical-biological functions and (Q) SARs. *Bioorganical Medicine Chemistry*, 15(6), 2223-2268. doi: 10.1016/j.bmc.2007.01.011
- Vincenti, M. P., & Brinckerhoff, C. E. (2002). Transcriptional regulation of collagenase (MMP-1, MMP-13) genes in arthritis: integration of complex signaling pathways for the recruitment of gene-specific transcription factors. *Arthritis Research & Therapy*, 4(3), 157-164. doi: 10.1186/ar401
- Visse, R., & Nagase, H. (2003). Matrix metalloproteinases and tissue inhibitors of metalloproteinases: Structure, function, and biochemistry. *Circulation Research*, 92(8), 827-839. doi: 10.1161/01.RES.000.0070112.80711.3D
- Wang, M., Sampson, E. R., Jin, H., Li, J., Ke, Q. H., Im, H., & Chen, D. (2013). MMP13 is a critical target gene during the progression of osteoarthritis. *Arthritis Research & Therapy*, 15(5), 1-11. doi: 10.1186/ar4133

- Woessner, J. F. (1991). Matrix metalloproteinases and their inhibitors in connective tissue remodeling. *The FASEB Journal*, 5(8), 2145-2154. doi: 10.1096 / fasebj.5.8.1850705
- Ye, S., Patodi, N., Walker-Bone, K., Reading, I., Cooper, C., & Dennison, E. (2007). Variation in the matrix metalloproteinase-3, -7, -12 and -13 genes is associated with functional status in rheumatoid arthritis. *International Journal of Immunogenetics*, 34(2), 81-85. doi: 10.1111 / j.1744-313X.2007.00664.x
- Yo, T. H., & Werb, Z. (1998). *Gelatinase B: structure, regulation, and function. Matrix metalloproteinases* (pp. 115-149). San Diego, CA: Academic Press.