

## Recovery of *Mollicutes* from the reproductive tract of dairy cattle in the state of Pernambuco, Brazil<sup>1</sup>

Sandra B. Santos<sup>2\*</sup>, José W. Pinheiro-Júnior<sup>3</sup>, André R. Mota<sup>2</sup>, André S. Santos<sup>2</sup>, Bruno H.L.S. Alves<sup>2</sup>, Júnior M.B. Oliveira<sup>3</sup>, Leonildo B.G. Silva<sup>2</sup> and Rinaldo A. Mota<sup>2</sup>

**ABSTRACT.-** Santos S.B., Pinheiro-Júnior J.W., Mota A.R., Santos A.S., Alves B.H.L.S., Oliveira J.M.B., Silva L.B.G. & Mota R.A. 2015. **Recovery of *Mollicutes* from the reproductive tract of dairy cattle in the state of Pernambuco, Brazil.** *Pesquisa Veterinária Brasileira* 35(6):491-496. Laboratório de Bacterioses dos Animais Domésticos, Departamento de Medicina Veterinária, Universidade Federal Rural de Pernambuco, Av. Dom Manoel de Medeiros s/n, Recife, PE 52171-900, Brazil. E-mail: [sanbsantos@gmail.com](mailto:sanbsantos@gmail.com)

The aim of the present study was to report the occurrence of members of the *Mollicutes* class in the reproductive system of dairy cattle in Brazil. Five farms containing dairy cattle were visited in January of 2012. In total, 100 cows of different ages, breeds and stages of lactation were examined in the present study. The cows were part of intensive or semi-intensive management systems and were submitted to mechanical milking or hand milking. The samples were collected after washing the vulvar region with water and soap, and then drying it with paper towels and disinfecting the area with alcohol (70°GL). Vaginal mucous was collected using a sterile alginate cotton swab, which was rubbed on the vagina, as well as the lateral and internal walls. Vulvovaginal mucous samples were cultured in both liquid and solid modified Hayflick's medium, for mycoplasmas, and UB medium, for ureaplasmas. The PCR assays for *Mollicutes* and *Ureaplasma* spp. were performed according to the standard protocols described in the current literature. During isolation, the frequency of *Mycoplasma* spp. was of 13.0% (13/100) and for *Ureaplasma* spp. was of 6.0% (6/100). In the PCR assays the frequency of *Mollicutes* was of 26.0% (26/100) and for *Ureaplasma* spp. was of 13.0% (13/100) in the dairy cattle studied. This is the first report of these agents in reproductive system of bovine of the Pernambuco state. Further studies are necessary to determine the pathogenic potential and species of these field isolates.

INDEX TERMS: Reproduction diseases, mycoplasmosis, dairy cows, vaginal mucous, *Ureaplasma* spp.

### RESUMO.- [Recuperação de *Mollicutes* do trato reprodutivo de bovinos leiteiros no Estado de Pernambuco.]

O presente estudo relata a ocorrência de membros da Classe *Mollicutes* no sistema reprodutivo de bovinos leiteiros no Brasil. Foram visitadas em janeiro de 2012 cinco fazendas de bovinos leiteiros. Um total de 100 vacas de diferentes idades, raças e estágios de lactação foram examinadas. Os animais foram mantidos em sistema de manejo inten-

sivo e/ou semi-intensivo, sendo submetidos aos sistemas de ordenha manual ou mecânica. As amostras de muco foram colhidas após a lavagem da região vulvar com água e sabão, com posterior desinfecção com álcool (70°GL). O muco vaginal foi colhido com suabe alginado estéril que foi friccionado nas paredes internas da vagina. Em seguida, as amostras foram cultivadas em meio Hayflick's modificado, para micoplasmas, e em meio UB, para ureaplasmas, ambos caldo e placa. Os ensaios de PCR para *Mollicutes* e *Ureaplasma* spp. foram realizados de acordo com protocolo padrão descrito na literatura. No isolamento, a frequência de *Mycoplasma* spp. foi de 13% (13/100) e para *Ureaplasma* spp. foi de 6% (6/100). Nas reações de PCR a frequência para *Mollicutes* foi de 26% (26/100) e para *Ureaplasmas* spp. foi de 13% (13/100) nos rebanhos bovinos leiteiros estudados. Este é o primeiro relato destes agentes no trato

<sup>1</sup> Received on January 17, 2015.

Accepted for publication on May 6, 2015.

<sup>2</sup> Laboratório de Bacterioses dos Animais Domésticos, Departamento de Medicina Veterinária, Universidade Federal Rural de Pernambuco (UFRPE), Av. Dom Manoel de Medeiros s/n, Recife, PE 52171-900, Brazil. \*Corresponding author: [sanbsantos@gmail.com](mailto:sanbsantos@gmail.com)

<sup>3</sup> Unidade Acadêmica de Garanhuns, UFRPE, Av. Bom Pastor s/n, Boa Vista, Garanhuns, PE 55292-270, Brazil.

reprodutivo de bovinos no Estado de Pernambuco. Estudos adicionais são necessários para determinar as espécies e o potencial patogênico destes isolados de campo.

TERMOS DE INDEXAÇÃO: Doenças da reprodução, micoplasmoses, vacas leiteiras, muco vaginal, *Ureaplasma* spp.

## INTRODUCTION

*Mollicutes* are bacteria that differ from other prokaryotes due to the lack of a cell wall and their genome size, which is the smallest reported (Herrmann 1992, Dybvig & Voelker 1996). These bacteria inflict a wide range of diseases on livestock and are generally associated with clinical manifestations such as pneumonia, conjunctivitis, polyarthritis, mastitis, agalactia, abortion and infertility (Nicholas 2002, McAuliffe et al. 2005). In cattle, the species of most relevance are in the group known as the *Mycoplasma mycoides* Cluster (MMC): *M. mycoides* subsp. *mycoides* Small Colony (*MmmSC*); the agent of contagious bovine pleuropneumonia (CBPP); and other mycoplasmas species such as *Mycoplasma bovis*, *Mycoplasma canadense*, *Mycoplasma arginini*, *Mycoplasma alkalescens*, *Mycoplasma bovirhinis*, *Mycoplasma bovigenitalium*, *Mycoplasma conjunctivae*, *M. mycoides mycoides capri*, as well as *Ureaplasma diversum* and *Acholeplasma laidlawii* (Landford 1975, Mulira et al. 1992, Razin et al. 1998, Frey 2002, Tenk 2005, Buzinhani et al. 2007, Marques et al. 2009, Santos et al. 2009). Some of these species are pathogenic, whereas others are considered ubiquitous of the normal flora in the mammary gland, as well as the respiratory and reproductive tracts (Whitford et al. 1994, Razin et al. 1998).

*Mycoplasma* spp. has often been associated with arthritis, pneumonia and otitis in neonatal calves (Yeruham et al. 1999, Gagea et al. 2006). *M. bovis* is one of the most economically important species of *Mycoplasma* in cattle herds, causing outbreaks of otitis, pneumonia and arthritis in calves, as well as highly infectious mastitis in cattle (Ghadersohi et al. 1999, Yeruham et al. 1999, Maeda et al. 2003, Francoz et al. 2004, Lamm et al. 2004, Radaelli et al. 2011). Economic losses caused by the syndrome (otitis-pneumonia-arthritis) in calves, as well as mastitis-agalactia in dairy cattle, have been previously studied in the Europe, United Kingdom, USA and Canada (Jasper 1981, Woldehiwet et al. 1990, Walz et al. 1997, Tenk 2005, Foster et al. 2007, Maunsell & Donovan 2009) and in the Brazil (Mettifogo

et al. 1996, Pretto et al. 2001, Nascimento et al. 2005). *M. bovirhinis* and *M. alkalescens* have also been isolated from cattle with mastitis (Jasper 1979, Jasper et al. 1979, Jasper 1981). Infected milk has been the source of *Mycoplasma* spp. in calves causing otitis and pneumonia. This agent is more prevalent in calves from herds with *Mycoplasma* mastitis than in calves from herds without *Mycoplasma* mastitis, and is considered a risk factor for outbreak infections in suckling calves (Jasper et al. 1979, Pftzner 1990, Lamm et al. 2004).

*Ureaplasma diversum* colonizes the urogenital region, where the infection has been associated with different clinical manifestations. In cattle, the main disorders are vulvitis, granular vulvovaginitis, endometritis, salpingitis, placentitis, fetal alveolitis, abortions and the birth of weak calves. In bulls, the main disorders are the presence of seminal vesiculitis, epididymitis, balanoposthitis and alterations in spermatozooids (Panangala et al. 1981, Miller et al. 1983, Ruhnke et al. 1984, Pilaszek & Truszczynski 1988, Cardoso et al. 2000, Nascimento et al. 2005, Oliveira Filho et al. 2005, Buzinhani et al. 2007, Rizzo et al. 2011). The agent is transmitted through organic secretions (semen, preputial and vaginal mucus) and direct contact during sexual intercourse, artificial insemination or embryo transfer (Kirkbride 1987, Hasso et al. 1993, Marques et al. 2009). In addition to reproductive diseases, *Ureaplasma* spp. has been reported in cases of destructive polyarthropathy (fetus) and mastitis in cattle, although these cases are rare (Himsworth et al. 2009). The aim of the present study was to describe the occurrence of agents of the *Mollicutes* Class in dairy cattle from different geographical areas in northeastern Brazil.

## MATERIALS AND METHODS

**Geographic area.** Five dairy cattle farms in the meso-region of the Agreste in the state of Pernambuco were visited in January of 2012. This geographical area is notable in the northeast of the country for its high milk production. The farms investigated were chosen at random in different cities, according to the following geographic coordinates: Garanhuns, municipal districts (Garanhuns: L 0773162 and UTM 9019050; Jupi: L 0565436 and UTM 9056244; Jurema: L 0813401 and UTM 9033624), Vale do Ipanema (Águas Belas: L 0716507 and UTM 8990870) Vale do Ipojuca (Alagoinha: L 0739156 and UTM 9059012), giving a total of five municipal districts (Fig.1).




Fig.1. Geographic area of the investigation for the presence of *Mollicutes* in dairy cattle in Brazil.

**Cows and sampling reproductive tract.** In total, 100 cattle of different ages, breeds and stages of lactation were examined in the present study. The cattle were part of intensive or semi-intensive management systems and were submitted to mechanical milking or hand milking. The sampling was performed using a stratified sampling farm, as described by Pereira (2003). Twenty, 15, 30, 23 and 20 cattle were randomly chosen from farms A, B, C, D and E, respectively.

The cattle were examined post milking. The vulvar and vaginal region of each animal was inspected for the presence of mucopurulent and purulent vaginal secretions or lesions. The medical history of the cattle herd was investigated for detected reproductive failure, endometritis, repeat breeding, infertility, the birth of weak calves and abortions. The samples were collected after washing the vulvar region with water and soap, and then drying it with paper towels and disinfecting the area with alcohol (70%GL). Vaginal mucous was collected using a sterile alginate cotton swab, which was rubbed on the vagina, as well as the lateral and internal walls. The samples were collected in sterile vials which were labeled with the number of the animal and farm. The samples were dipped into transport solution (phosphate buffered saline, pH 7.2), maintained at 4°C in a cooler with ice and taken to the Laboratory of Infectious Diseases (LDIC-DMV/UFRPE).

**Bacteriological procedures.** The vulvovaginal mucous samples were cultured for mycoplasmas and ureaplasmas, as described by Razin & Tully (1996) and Ruhnke & Rosendal (1994). Previously, 2mL of each sample were filtered through a sterile syringe holder coupled with a millipore membrane (0.45µm diameter). Then, 100µL of the filtrate was diluted up to 10<sup>-1</sup> at 10<sup>-5</sup> and inoculated in both liquid and solid modified Hayflick's medium, that optimizes the isolation of *Mycoplasma* classic and UB medium, selective for the propagation of ureaplasmas. Subsequently, all samples sowed were incubated at 37°C for a minimum period of up to 21 days. The plates were placed in a jar to determine the microaerophilic conditions. The agar plates were checked daily for the presence of colonies using a stereomicroscope (40X). Blind sub cultures were used for up to 60 days, before disposal of the samples. *Mollicutes* isolates were confirmed by Dienes probe; *Mycoplasma* genera was used a digitonin sensitivity test and *Ureaplasma* spp. colonies were identified by the urease production in the U4 medium and their characteristic colony morphology (Whitford et al. 1994).

**DNA extraction and PCR assays.** The DNA was extracted from the samples by the method described in the commercially available kit (DNA Easy Blood and Tissues Kit®, Qiagen Biotechnology, guideline page 25). The primers (GPO-3 and MGSO) that amplify the V6 and V7 conserved regions of the 16S RNA gene, specific to *Mollicutes* class was used as a procedure for screening of field samples (Van Kuppeveld et al. 1992). The samples were also submitted a PCR assays to detect *Ureaplasma* spp. using primers UGP-F' (gene location 257-256) and UGP-R' (gene location 862-881) conserved regions of the 16S RNA gene for *Ureaplasma* spp. (Lauerma 1998). The PCR assays for *Mollicutes* were performed in a mixture prepared with a volume of 25µL, containing 5µL of DNA template, 30pmol of each primer, MgCl<sub>2</sub> (1.5mM), buffer [10mM of Tris-HCl, pH 8.3], a mix of deoxynucleoside triphosphate (50µM), *Taq* DNA Polymerase (2.5U) and Milli-Q ultrapure water. For *Ureaplasma* spp., PCR assays were performed with primers at 30pmol, Milli-Q ultrapure water and 6.25µL of TopTaq Mastermix, following the manufacturer's instructions. In vitro amplifications were performed using thermo cycler model PTC-100 (MJ-Research®). Ultrapure water was used as a negative control. *Mycoplasma mycoides mycoides* (reference strain GM12) and *Ureaplasma diversum* (reference strain GMU132, Vup6) were

used as positive controls in each reaction, respectively. The amplified PCR was visualized by electrophoresis in 1.5% agarose gel with 100bp molecular weight marker, colored with Bluegreen, viewed under ultraviolet light and photodocumented.

## RESULTS

On the farms studied, the cattle examined were asymptomatic for mycoplasmosis and other reproductive disorders during the period of the research. In the swabs from their reproductive systems, the total prevalence for *Mollicutes* was 19.0% (19/100) in the isolation and 39.0% (39/100) in the PCR. In the samples cultivated in agar Hayflick's medium, 13.0% growth was recorded and it was possible to visualize flask-shaped cells, which in the digitonin test was confirmed as *Mycoplasma* spp. In the agar UB medium, 6.0% of the samples exhibited characteristics of *Ureaplasma* spp. typical colonial morphology of brown gold colonies, confirmed by urease test. The subculture was made for growth confirmation of *Mycoplasma* spp. and *Ureaplasma* spp. The PCR confirmed 26.0% for *Mollicutes* the amplicon obtained was of 270bp and 13.0% for *Ureaplasma* spp. with amplicon of 644bp (Fig.2).

## DISCUSSION

Mycoplasmas are ubiquitous in nature and frequently contain microbiota from different systems, such as the ear, eye, joints and mammary gland, as well as the respiratory and genitourinary tract in animals, humans and insects (Baseman & Tully 1997, Dybvig & Voelker 1996). The findings of the present study confirmed that *Mycoplasma* spp. e *Ureaplasma* spp. strains grew on both solid and liquid media and were confirmed at the genera level by the PCR. The presence of *U. diversum* in the reproductive tract of cattle with or without symptoms of mycoplasmosis has been reported in the literature by several authors (Oliveira Filho et al. 2005, Buzinhani et al. 2007, Razin et al. 1998). These agents in the reproductive tract of cattle may or may not be associated with outbreaks of reproductive disorders. Ho-




Fig.2. Results of the PCR for *Ureaplasma* spp. in the vaginal mucous of dairy cows, state of Pernambuco, Brazil. Lane M, molecular size marker (Amresco® 1kb, amplisize standard 500-10,000); Lanes 1,2,4,5,7,9,11 DNA *Ureaplasma* spp.; Lane (C-), negative control; Lanes (C+) Positive control.



wever, the genus *Ureaplasma* spp. has often been associated with disease outbreaks (Albertsen 1955, Onoviran et al. 1975, Britton et al. 1987, Le Grand et al. 1995, Cardoso et al. 1997, Nascimento et al. 1998, Cardoso et al. 2000, Cardoso & Vasconcelos 2004, Nascimento et al. 2005). In cattle, it can cause placentitis, fetal alveolitis, granular vulvovaginitis, abortion and the birth of weak calves and infection. In bulls, it may result in seminal vesiculitis, balanoposthitis and alterations in spermatozooids (Howard et al. 1976, Mulira & Saunder 1994, Himsworth et al. 2009, Marques et al. 2009). The occurrence of *Ureaplasma* spp. in cattle could be a potential risk for manifestations of reproductive diseases that have already been reported in the literature. The possible transmission route is through direct contact during sexual intercourse or by artificial insemination. This bacteria is shed in semen, milk, preputial and vaginal mucus (Kirkbride 1987, Britton et al. 1988, Cardoso & Vasconcelos 2004).

In Brazil, there are very few studies reporting *Ureaplasma* infection in cattle herds. However, Nascimento et al. (1998) isolated *M. bovirhinis* in the vaginal mucus of heifers, highlighting the importance of this species as a cause of reproductive failure in this country, as well as a probable decrease in milk and meat production, similar to what occurs in others countries. *M. bovis* was diagnosed in 57% of the samples of vaginal mucus from cattle with reproductive disorders (vulvovaginitis, abortion, stillborn) and the authors highlighted the risk of venereal transmission of *Mycoplasma* spp. (Nascimento et al. 2005). In the northeast of Brazil, Santos et al. (2013) reported *Mollicutes* with frequencies of 65.6% and for *Ureaplasma diversum* of 15.6% in reproductive disease outbreak in cattle in Paraíba, the cows presented heat repetition, granular vulvovaginitis and abortions.

In the present study, the occurrence of *Mycoplasma* and *Ureaplasma* spp. in the vaginal mucus was low in comparison to the frequencies reported in other Brazilian states (Cardoso et al. 1997, Cardoso et al. 2000, Cardoso & Vasconcelos 2004, Buzinhani et al. 2007, Santos et al. 2013). However, it was possible to prove the viability of these microorganisms through bacterial isolation, thus highlighting the possibility of these agents causing reproductive diseases in these cattle during opportunistic outbreaks. Further studies are required to prove the species and pathogenicity of these isolates. In the cattle assessed in the present study, there was no history of reproductive disorders or abortion, although the presence of these agents in the vaginal mucus of cows could be considered a risk factor for reproductive disease.

## CONCLUSION

It was possible to verify the occurrence of *Mycoplasma* spp. and *Ureaplasma* spp. in dairy cattle from different geographical areas of the state of Pernambuco. This is the first report of these bacteria in this region of Brazil. Further studies are necessary to determine the species and pathogenic potential of these field isolates.

**Acknowledgements.**- This study was supported by the Conselho Nacional de Desenvolvimento Científico e Tecnológico - CNPq (MCT/CNPq) and

Fundação de Amparo à Ciência e Tecnologia do Estado de Pernambuco - FACEPE (Programa de Desenvolvimento Científico Regional - DCR nº 0006-5.05/10 and APQ 1512-5.05/10). To Dr. Elmiro Rosendo do Nascimento by providing standards strains used on the project.

**Conflicts of interest.**- The authors have no conflicts of interest to declare.

**Research Ethics Committee.**- This project was approved by the ethics committee of the UFRPE/DMV Process nº 23082.004146/2011-28.

## REFERENCES

- Albertsen B.E. 1955. Pleuropneumonia-Like organisms in the semen of Danish artificial insemination bulls. *Nordish Veterinaer Medicin* 7:169-201.
- Baseman J.B. & Tully J.G. 1997. Mycoplasmas: sophisticated, reemerging, and burdened by their notoriety. *Emerg. Infect. Dis.* 3:21-32.
- Britton A.P., Ruhnke H.L., Miller R.B., Johnson W.H., Leslie K.E. & Rosedal S. 1987. In vitro exposure of bovine morulae to *Ureaplasma diversum*. *Can. J. Vet. Res.* 51:198-203.
- Britton A.P., Miller R.B., Ruhnke H.L. & Johnson W.H. 1988. The recovery of ureaplasmas from bovine embryos following in vitro exposure and ten washes. *Theriogenology* 30:997-1003.
- Buzinhani M., Metiffogo E. & Timenetsky J. 2007. Detecção de *Mycoplasma* spp. e *Ureaplasma diversum* em vacas com distúrbios reprodutivos. *Arq. Bras. Med. Vet. Zootec.* 59:1368-1375.
- Cardoso M.V. & Vasconcelos S.A. 2004. Importância das micoplasmoses na infertilidade de touros. *Arqs Inst. Biológico, São Paulo*, 71:257-265.
- Cardoso M.V., Scarcelli E., Grasso L.M.P.S., Teixeira S.R. & Genovez M.E. 2000. *Ureaplasma diversum* and reproductive disorder in Brazilian cows and heifers, first report. *Anim. Reprod. Sci.* 63:137-143.
- Cardoso M.V., Grasso L., Stefano E., Okuda L.H. & Cunha R.A.F. 1997. Isolamento de *Ureaplasma diversum* e *Mycoplasma* spp. em casos de Vulvite Granular Bovina. *Revta Bras. Reprod. Anim.* 21:172-173.
- Dybvig K. & Voelker L.L. 1996. Molecular biology of mycoplasmas. *Annu. Rev. Microbiol.* 50:25-57.
- Foster A.P., Naylor R.D., Howie N.M., Nicholas R.A.J. & Ayling R.D. 2007. *Mycoplasma bovis* and otitis in dairy calves in the United Kingdom. *Vet. J.* 179:455-457.
- Francoz D., Fecteau G., Desrochers A. & Fortin M. 2004. Otite media in dairy calves: A retrospective study of 15 cases (1987 to 2002). *Can. Vet. J.* 45:661-666.
- Frey J. 2002. Mycoplasmas of animals, p.73-90. In: Razin S. & Herrmann R. (Eds), *Molecular Biology and Pathogenicity of Mycoplasmas*. Kluwer Academic, New York.
- Gagea M.I., Bateman K.G., Shanahan R.A., Van Dreumel T., McEwen B.J.S., Archambault M. & Caswell J.L. 2006. Naturally occurring *Mycoplasma bovis*-associated pneumonia and polyarthritis in feedlot beef calves. *J. Vet. Diagn. Invest.* 18:29-40.
- Ghadersohi A., Hirst R.G., Forbes-Faulkener J. & Coelen R.J. 1999. Preliminary studies on the prevalence of *Mycoplasma bovis* mastitis in dairy cattle in Australia. *Vet. Microbiol.* 65:185-194.
- Hasso S.A., Al-Aubaidi J.M. & Al-Darraj A.M. 1993. Contagious agalactia in goats: it's severity as related to the route of infection and pregnancy. *Small Rumin. Res.* 10:263-275.
- Herrmann R. 1992. Genome Structure and Organization, p.157-68. In: Maniloff J., McElhaney R.N., Finch L.R. & Baseman J.B. (Eds), *Mycoplasmas: molecular biology and pathogenesis*. Am. Soc. Microbiol., Washington, DC.
- Himsworth C.G., Hill J.E., Huang Y., Waters E.H. & Wobeser G.A. 2009. Destructive polyarthropathy in aborted bovine fetuses: a possible association with *Ureaplasma diversum* Infection? *Vet. Pathol.* 46:269-272.
- Howard C.J., Gourlay R.N., Thomas L.H. & Stott E.J. 1976. Induction of pneumonia in gnotobiotic calves following inoculation of *Mycoplasma disparand* ureaplasmas (T-mycoplasmas). *Res. Vet. Sci.* 21:227-231.
- Jasper D. 1979. Bovine mycoplasmal mastitis. *J. Am. Vet. Med. Assoc.* 175: 1072-1074.

- Jasper D. 1981. Bovine mycoplasmal mastitis. *Adv. Vet. Sci. Comp. Med.* 25:121-159.
- Jasper D., Dellinger J. & Rollins M. 1979. Prevalence of *Mycoplasma* bovine mastitis in California. *Am. J. Vet. Res.* 40:1043-1047.
- Kirkbride C.A. 1987. *Mycoplasma*, *Ureaplasma*, and *Acholeplasma* infections of bovine genitalia. *Vet. Clin. North Am., Food Anim. Pract.* 3:575-591.
- Lamm C.G., Munson L., Thurmond M.C., Barr B.C. & George L.W. 2004. *Mycoplasma* otitis in California calves. *J. Vet. Diagn. Invest.* 16:397-402.
- Landford E.V. 1975. *Mycoplasma* species recovered from the reproductive tracts of Western Canadian cows. *Can. J. Comp. Med.* 39:133-137.
- Lauerman L.H. 1998. Mycoplasma PCR assays, p.41-42. In: Lauerman L.H. (Ed.), *Nucleic Acid Amplification Assays for Diagnosis of Animal Diseases*. American Association of Veterinary Laboratory Diagnosticians, Turlock, CA.
- Le Grand D., Poumarat F. & Martel J.L. 1995. Genitale infection by *Ureaplasma diversum*: survey in cattle in France. *Vet. Res.* 26:11-20.
- Maeda T., Shibahara T., Kimura K., Wada Y., Sato K., Imada Y., Ishikawa Y. & Kadota K. 2003. *Mycoplasma bovis* associated suppurative otitis media and pneumonia in Bull calves. *J. Comp. Pathol.* 129:100-110.
- Marques L.M., Buziniani M., Neto R.L., Oliveira R.C., Yamaguti M., Guimarães A.M. & Timenetsky J. 2009. Detection of *Ureaplasma diversum* in bovine semen straws for artificial insemination. *Vet. Rec.* 165:572-573.
- Maunsell F.P. & Donovan G.A. 2009. *Mycoplasma bovis* Infections in young calves. *Vet. Clin. Food Anim.* 25:139-177.
- McAuliffe L., Ellis R.J., Lawes J.R., Ayling R.D. & Nicholas R.A.J. 2005. 16S rDNA PCR and denaturing gradient gel electrophoresis: a single generic test for detecting and differentiating *Mycoplasma* species. *J. Med. Microbiol.* 54:731-739.
- Mettifogo E., Nascimento E.R., Muller E.E., Nascimento M.G.F. & Freitas J.C. 1996. Mastite bovina por *Mycoplasma bovis*. *Revta Bras. Med. Vet.* 18:22-25.
- Miller R.B., Ruhnke H.L., Doig P.A., Poitras B.J. & Palmer N.C. 1983. The effects of *Ureaplasma diversum* inoculated into the amniotic cavity in cows. *Theriogenology* 20:367-374.
- Mulira G.L. & Saunders J.R. 1994. Humoral and secretory antibodies to *Ureaplasma diversum* in heifers following subcutaneous vaccination and vaginal infection. *Can. J. Vet. Res.* 58:104-108.
- Mulira G.L., Saunders R.J. & Barth D.A. 1992. Isolation of *Ureaplasma diversum* and mycoplasma from genital tracts of beef and dairy cattle in Saskatchewan. *Can. Vet. J.* 33:46-49.
- Nascimento M.G.F., D'Angelis F.H.F., Nascimento E.R., Resende O.A. & Lignon G.B. 1998. *Mycoplasma bovirhinis* em muco vaginal de novilhas. *Revta Bras. Med. Vet.* 20:210-214.
- Nascimento M.G.F., D'Angelis F.H.F., Nascimento E.R. & Resende O.A. 2005. Envolvimento de micoplasmas em vacas com distúrbios reprodutivos. *Acta Scient. Vet.* 33:195-199.
- Nicholas R.A.J. 2002. Improvements in the diagnosis and control of diseases of small ruminants caused by mycoplasmas. *Small Rumin. Res.* 45:145-149.
- Oliveira Filho B.D., Porto R.N.G., Gambarini M.L., Kunz T.L., Ferraz H.T., Viu M.A.O., Lopes D.T. & Sousa A.P.F. 2005. Isolamento de *Ureaplasma diversum* em muco vulvovaginal de vacas leiteiras repetidoras de estro no Estado de Alagoas, Brasil. *Arch. Vet. Sci.* 10:151-156.
- Onoviran O., Truscott R.B., Fish N.A., Barker C.A.V. & Ruhnke H.L. 1975. The recovery of mycoplasmas from the genital tracts of bulls in artificial breeding units in Ontario. *Can. J. Comp. Med.* 39:474-475.
- Panangala V.S., Winter A.J., Wijesinha A. & Foote R.H. 1981. Decreased motility of bull spermatozoa caused by *Mycoplasma bovirhinis*. *Am. J. Vet. Res.* 42:2090-2093.
- Pereira M.G. 2003. *Epidemiologia Teoria e Prática*. 7th ed. Guanabara Koogan, Rio de Janeiro. 596p.
- Pfutzner H. 1990. Epizootiology of the *Mycoplasma bovis* infection of cattle. *Zentralbl. Bakteriol.* 20(Suppl.):394-399.
- Pilaszek J. & Truszczynski M. 1988. Affinity of microorganisms of the genus *Ureaplasma* to the reproductive organs of cattle. *Comp. Immunol. Microbiol.* 11:177-180.
- Pretto L.G., Muller E.E., Freitas J.C., Mettifogo E., Buziniani M., Yamaguti M. & Salvador R. 2001. Mastite bovina por *Mycoplasma bovis* em rebanhos leiteiros. *Pesq. Vet. Bras.* 21:143-145.
- Radaelli E., Castiglioni V., Losa M., Benedetti V., Piccinini R., Nicholas R.A.J., Scanziani E. & Luini M. 2011. Outbreak of bovine clinical mastitis caused by *Mycoplasma bovis* in a North Italian herd. *Res. Vet. Sci.* 91:251-253.
- Razin S. & Tully J.G. 1996. *Molecular and Diagnostic Procedures in Mycoplasma*. 2nd ed. Academic Press, California. 463p.
- Razin S., Yogeve D. & Naot Y. 1998. Molecular biology and pathogenicity of mycoplasmas. *Microbiol. Mol. Biol. Res.* 62:1094-1156.
- Rizzo H., Meira Junior E.B.S., Oliveira R.C., Yamaguti M., Buziniani M., Timenetsky J. & Gregory L. 2011. Isolamento e PCR para detecção de *Mollicutes* em muco vaginal e sua associação com problemas reprodutivos em ovinos criados na região de Piedade, São Paulo, Brasil. *Ciência Rural* 41:324-329.
- Ruhnke H.L. & Rosendal S. 1994. Useful protocols for diagnosis of animal mycoplasmas, p.141-155. In: Whitford H.W., Rosenbusch R.F. & Lauerman L. (Eds), *Mycoplasmosis in Animals: Laboratory Diagnosis*. Iowa State University Press, Ames, IA.
- Ruhnke H.L., Palmer N.C., Doig P.A. & Miller R.B. 1984. Bovine abortion and neonatal death associated with *Ureaplasma diversum*. *Theriogenology* 21:295-301.
- Santos S.B., Nascimento E.R., Faccini J.L.H., Barreto M.L. & Pereira V.L.A. 2009. Potentially pathogenic mycoplasmas in the external ear canal of clinically normal cattle in southeast Brazil: first report. *Braz. J. Microbiol.* 40:455-457.
- Santos S.B., Pinheiro-Júnior J.W., Oliveira A.A.F., Mota A.R., Oliveira J.M.B., Veras G.A., Nascimento E.R. & Mota R.A. 2013. Ocorrência de *Mollicutes* e *Ureaplasma* spp. em surto de doença reprodutiva em rebanho bovino no Estado da Paraíba. *Pesq. Vet. Bras.* 33:315-318.
- Tenk M. 2005. Examination of *Mycoplasma bovis* infection in cattle. Doctoral Thesis, Szent István University, Postgraduate School of Veterinary Science, Budapest. 70p.
- Van Kuppeveld F.J.M., Van Der Logt J.T.M., Angulo A.F., Zoest M.J.V., Quint W.G.V., Niesters H.G.M., Galama J.M.D. & Melchers W.J.G. 1992. Genus and species specific identification of mycoplasmas by 16S rRNA amplification. *Appl. Environ. Microbiol.* 58:2606-2615.
- Walz P., Mullaney T. & Render J.A. 1997. Otitis media in preweaned Holstein dairy calves in Michigan due to *Mycoplasma bovis*. *J. Vet. Diagn. Invest.* 9:250-254.
- Whitford H.W., Rosenbusch R.F. & Lauerman L.H. 1994. *Mycoplasmosis in Animals: laboratory diagnosis*. Iowa State University Press, Ames, IA. 173p.
- Woldehiwet Z., Mamache B. & Rowan T.G. 1990. Effects of age, environmental temperature and relative humidity on the colonization of the nose and trachea of calves by *Mycoplasma* spp. *Brit. Vet. J.* 146:419-424.
- Yeruham I., Elad D. & Liberbol M. 1999. Clinical and microbiological study of an otitis media outbreak in calves in a dairy herd. *J. Vet. Med. B* 46:145-150.