

Frequency, serotyping and antimicrobial resistance pattern of *Salmonella* from feces and lymph nodes of pigs¹

João B.P. Guerra Filho², Ricardo S. Yamatogi², Fábio S. Possebon², Sueli A. Fernandes³, Monique R. Tiba-Casas³, Gustavo H.B. Lara², Márcio G. Ribeiro² and José P.A.N. Pinto^{2*}

ABSTRACT.- Guerra Filho J.B.P., Yamatogi R.S., Possebon F.S., Fernandes S.A., Tiba-Casas M.R., Lara G.H.B., Ribeiro M.G. & Pinto J.P.A.N. 2016. **Frequency, serotyping and antimicrobial resistance pattern of *Salmonella* from feces and lymph nodes of pigs.** *Pesquisa Veterinária Brasileira* 36(12):1165-1170. Departamento de Higiene Veterinária e Saúde Pública, Faculdade de Medicina Veterinária e Zootecnia Universidade Estadual Paulista, Distrito de Rubião Júnior s/n, Botucatu, SP 18618-970, Brazil. E-mail: josepaes@fmvz.unesp.br

Salmonellosis is a foodborne disease caused by bacteria of the genus *Salmonella*, being pigs and pork-products potentially important for its occurrence. In recent decades, some serovars of *Salmonella* have shown increase of resistance to conventional antimicrobials used in human and animal therapy, with serious risks for public health. The aim of this study was to evaluate feces (n=50), mediastinal (n=50), mesenteric (n=50) and mandibular (n=50) lymph nodes obtained from slaughter houses for *Salmonella* spp. Positive samples were serotyped and subjected to an *in vitro* antimicrobial susceptibility test, including the extended-spectrum beta-lactamase (ESBL) production. *Salmonella* species were identified in 10% (20/200) of total samples. From these, 20% (10/50) were identified in the submandibular lymph nodes, 18% (9/50) in the mesenteric lymph nodes, 2% (1/50) in feces and 0% (0/50) in the mediastinal lymph nodes. The serotypes found were *Salmonella* Typhimurium (55%), *S. enterica* subsp. *enterica* 4,5,12: i: - (35%), *S. Brandenburg* and *S. Derby* with 5% (5% each). All strains showed resistance to at least one antimicrobial; 90% were resistant to four or more antimicrobials, and 15% were multidrug-resistant. Resistance to ciprofloxacin, tetracycline and nalidixic acid was particularly prevalent amongst the tested serovars. Here, we highlighted the impact of pigs in the epidemiological chain of salmonellosis in domestic animals and humans, as well as the high antimicrobial resistance rates of *Salmonella* strains, reinforcing the necessity for responsible use of antimicrobials for animals as an emergent One Health issue, and to keep these drugs for human therapy approaches.

INDEX TERMS: Serotyping, antimicrobial resistance, *Salmonella*, feces, lymph nodes, pigs, pork, multidrug-resistant, epidemiology, serovars.

RESUMO. [Isolamento, sorotipagem e padrões de resistência a antimicrobianos de *Salmonella* em fezes e linfonodos de suínos.] Nas últimas décadas, o aumento de cepas circulante de *Salmonella* concomitantemente

a resistência microbiana tem despertado a preocupação dos órgãos de Saúde Pública. Deste modo, o objetivo do presente trabalho foi pesquisar a presença de *Salmonella* a partir de fezes (n=50), linfonodos mediastinos (n=50), mesentéricos (n=50) e submandibular (n=50) oriundos de um abatedouro suíno. As cepas isoladas foram sorotipadas e testadas quanto a resistência antimicrobiana. A presença de *Salmonella* isolada foram em 10% (20/200) do total de amostras, sendo 20% dos linfonodos submandibulares, 18% dos linfonodos mesentéricos e 2% das fezes. Os sorotipos encontrados foram *S. Typhimurium* (55%), *S. enterica* subsp. *enterica* 4,5,12: i: - (35%), *S. Brandenburg* (5%) e *S.*

¹ Received on November 20, 2015.

Accepted for publication on June 16, 2016.

² Departamento de Higiene Veterinária e Saúde Pública, Faculdade de Medicina Veterinária e Zootecnia, Universidade Estadual Paulista (Unesp), Distrito de Rubião Júnior s/n, Botucatu, SP 18618-970, Brazil. *Corresponding author: josepaes@fmvz.unesp.br

³ Instituto Adolfo Lutz, Av. Doutor Arnaldo 351, São Paulo, SP 01246-902, Brazil.

Derby (5%). Todas as cepas apresentaram resistência a pelo menos um antimicrobiano testado, sendo 90% resistente a pelo menos quatro antimicrobianos. Destes, 15% foram classificadas como multidrogas resistentes. Os antimicrobianos mais resistentes entre os sorovares isolados foram a ciprofloxacina, tetraciclina e o ácido nalidixico. A presença de cepas de *Salmonella* resistente a antimicrobianos na espécie suína tem gerado um grande impacto epidemiológico entre homem e animal, reforçando cada vez mais a necessidade do uso adequado de drogas principalmente relacionado com o tema "One Health".

TERMOS DE INDEXAÇÃO: Isolamento, sorotipagem, resistência, antimicrobianos, *Salmonella*, fezes, linfonodos, suínos, multidrogas resistentes, epidemiologia, sorovares.

INTRODUCTION

Foodborne diseases always offer great risks, despite the attention given for their control. The increased production in a global scale has contributed to foodborne cases and outbreaks, resulting on serious public health concern (Carrasco et al. 2012). Today, salmonellosis is the most frequent foodborne disease in the world (Bollaerts et al. 2008), reaching 131,468 annual human cases reported in the European Union (Carrasco et al. 2012), and 45,828 in the United State (CDC 2013), even when the occurrence of mild symptoms leads to an underestimated notification (Santos et al. 2002).

Brazil is considered the 4th largest producer of pork in the world, second only to China, European Union and United States, respectively, reaching an annual production over 3 million tons and exports over 600 thousand tons in 2013 (ABIPECS 2013). In this country, from 2000 to 2013, were reported 8,871 foodborne diseases outbreaks, being *Salmonella* the causative agent in 1,522 of them. Among these total outbreaks, 277 (4.26%) were associated with pork products.

Salmonella is closely related to the swine production chain, being isolated in several production steps, including primary production, transport of animals, the pre-slaughter and pre-evisceration steps, especially during scalding, opening of the abdomen and withdrawing of colon (Letellier et al. 2009, Duggan et al. 2010, Carascvo et al. 2012, Gomes-Neves et al. 2012). *Salmonella* infections in pigs occurs through oral infection, and later spreads to the lymphatic system, which acts as barrier at the first moment, but can become reservoirs posteriorly allowing environmental-elimination of the agent and its dissemination through other animals (Straw et al. 2012). Therefore, the isolation of *Salmonella* from pigs' lymph nodes indicates its carrier status and the analysis of intestinal contents is related to its excretory potential (Davies et al. 1998, Bahnson et al. 2006).

In recent decades, *S. Enteritidis* and *S. Typhimurium* serovars have caused great concern (Bollaerts et al. 2008). In most cases where humans have contracted one of these serovars, the pathogen infection is self-limiting, characterized by gastroenteritis. However, in some cases the manifestations are more severe, especially in immunocompromised patients (EFSA 2010). In these cases, the conventional treatment for salmonellosis is based on fluoroquinolones

and quinolones antimicrobials for adults, third-generation cephalosporins for children, and chloramphenicol in patients with endocarditis or endovascular infection (Lesser & Miller 2005, EFSA 2010). Nevertheless, several studies have pointed out the isolation of multidrug-resistant *Salmonella* strains, including to the main drugs of choice in the therapeutic practices for veterinary and human protocols (EFSA 2010).

In the European Union is recorded occurrence of multidrug-resistant strains in 16 countries for pigs, and 14 countries for pork samples (EFSA 2010). In Spain, studies report *Salmonella* resistance to streptomycin (46% of tested serovars), tetracycline (30%), sulfonamides (25%) and ampicillin (23%), with 36% of serovars multidrug-resistant (Gomez-Laguna et al. 2011). In a similar study conducted in Vietnam, from poultry and pork meat strains, resistance of *Salmonella* to at least one antimicrobial was found in 78.4% of samples, with 23.2% MDR (Thai et al. 2012). In a study carried out in Brazil, different serotypes of *Salmonella* isolated from pigs showed resistance to sulfonamides (83.9%), tetracycline (37.4%), cotrimoxazole (25.2%), ampicillin (20.2%), chloramphenicol (16.1%), streptomycin (14.1%), and nalidixic acid (10.1%), and 24.2% were formally multidrug-resistant (Castagna et al. 2001).

It is well recognized that the *Enterobacteriaceae* family includes many species that produce enzymes that hydrolyze beta lactam antibiotics. One of the predominant enzymes, Extended Spectrum Beta Lactamase (ESBL), inactivates penicillins, cephalosporins and monobactams (Souza Junior et al. 2004).

Here, we investigated the frequency of *Salmonella* spp. in feces, mesenteric, mediastinal and submandibular lymph nodes from pigs, as well as serotype characterization and *in vitro* resistance profile of strains to several antimicrobials, chosen based on Clinical Laboratory Standards Institute guidelines (CSLI 2013).

MATERIALS AND METHODS

Animals and sample collection. Two hundred specimens from pigs were sampled. Of these, 50 fecal samples were collected in the evisceration and inspection tables, using sterile plastic bags and 150 lymph nodes without apparent abnormalities (without lymphadenitis) of pigs were removed from carcasses and placed in sterile plastic bags (being 50 mediastinal, 50 mesenteric and 50 submandibular), taken randomly from different animals. The animals were slaughtered in the finishing phase (150-180 days) in slaughterhouses under Brazilian Federal Inspection Service. Sampled pigs came from up 20 piggeries of 10 different cities located in Sao Paulo and Santa Catarina States, Brazil. The animals were from medium scale farms housing between 350-950 animals kept in intensive indoors system, with concrete-floored, and fed exclusively with commercial feed. Immediately after collection, the samples were kept refrigerated (4-8°C) and stored at -20°C until the diagnostic procedures.

***Salmonella* identification.** For the isolation of *Salmonella* spp., the samples defrost under refrigeration for 24 hours. Feces samples were fractionated in 1g aliquots and packed in sterile plastic bags. Samples of lymph nodes were externally disinfected with alcohol 70% and then fractionated to obtain a 1g aliquot, which was also transferred to sterile plastic bag. For each bag was added 9mL of buffered peptone water 1% (BPW) (Oxoid®,

Hampshire - England), homogenized and incubated for 24h at 35 °C. Subsequently the enrichment step, were transferred 0.1mL of pre-enrichment to 10mL of Rappaport-Vassiliadis broth (RV) and 1mL to 10mL of Tetrathionate broth (TT). TT tubes were incubated at 35°C for 18 to 24 hours and RV incubated 42°C for 18 to 24 hours. Each broth was then streak on plates containing xylose lysine desoxycholate (XLD - Oxoid®), bismuth sulfite agar (BS - Oxoid®) and incubated for 24 hours at 35°C. The colonies compatible with *Salmonella* spp. were transferred to triple sugar iron agar (TSI - Oxoid®) and lysine iron agar (LIA - Oxoid®) for preliminary biochemical tests, incubated for 18 to 24 h at 35°C. *Salmonella* suspected colonies were subjected to follow conventional phenotypic tests: indole, motility, phenylalanine desaminase and urease production, Methyl Red and Voges-Proskauer reactions, citrate, glucose and lactose utilization. The samples with typical biochemical characteristics were confirmed by agglutination test in polyvalent antiserum specific for *Salmonella* spp (Probac®, São Paulo, Brazil).

Microbiological identification of *Salmonella* spp. was performed according to Andrews et al. (2014). The strains were serotyped using the somatic O, phase 1 and phase 2 of the H flagellar antigens by agglutination tests with antisera prepared in the Laboratory of Enteric Pathogens, Institute Adolfo Lutz, São Paulo as specified in the White-Kauffmann-LeMinor protocol for *Salmonella* serotyping (Grimont & Weil 2007).

In vitro antimicrobial susceptibility test. For the antimicrobial resistance investigation, the strains were subjected to the Bauer-Kirby disk diffusion method, according to the guidelines recommended by Clinical and Laboratory Standards Institute (CLSI, 2013). The antimicrobial agents were: nalidixic acid - 30 g (NAL), amikacin - 30µg (AMI), ampicillin - 10µg (AMP), aztreonam - 30µg (AZT), cefepime - 30µg (CPM), cefotaxime - 30µg (CTX), ceftazidime - 30µg (CAZ), ceftriaxone - 30µg (CRO), ciprofloxacin - 5µg (CIP), chloramphenicol - 30µg (CLO), streptomycin - 10µg (EST), gentamicin - 10µg (GEN), meropenem (MER), sulfonamides - 300µg (SUL), trimethoprim-sulfamethoxazole - 1.25/23.75µg (SUT), and tetracycline - 30µg (TET). In the current study, the intermediate resistant strains were considered resistant (CDC 2014).

ESBL production. Ciprofloxacin resistant strains were tested for ESBL enzyme presence by the method of double disk diffusion to check synergism between clavulanic acid and aztreonam, ceftazidime, ceftriaxone and cefotaxime. Resistant strains were identified by the formation of a secondary inhibition halo by a synergism of the tested antimicrobials with the initial intersection of the inhibition zones of each antimicrobial (CSLI 2013).

Statistical analysis. Logistic regression analysis was used in the binomial dependent variable, using the PROC LOGISTIC of the Statistical Analysis Software (SAS® 9.4, Inst. Inc., Cary, NC, USA). The results are shown as a percentage. For all analyzes, were adopted the significance level of 5% ($P < 0.05$).

RESULTS

Salmonella strains were identified in 10% ($n=20/200$) of samples. The highest rates were found in the submandibular lymph nodes ($n=10/50$, 20%), followed by mesenteric lymph nodes ($n=9/50$, 18%). In contrast, the lowest prevalence were found in fecal samples ($n=1/50$, 2%). No strain of the pathogen was isolated in mediastinal lymph nodes. Mesenteric and submandibular lymph nodes showed statistical similarity ($p > 0.05$) but differed when compared to samples of feces and mediastinal lymph nodes ($p < 0.05$) (Table 1).

The serotyping of the 20 isolated strains identified the following serotypes: *S. Typhimurium* ($n=11/20$, 55%), *S. enterica* subsp. *enterica* 4,5,12: i: - ($n=7/20$, 35%), and *S. Brandenburg* and *S. Derby* ($n=1/20$, 5% each).

The highest antimicrobial resistance rates of isolates were found for ciprofloxacin (CIP) and tetracycline (TET) ($n=18/20$, 90% each), followed by nalidixic acid (NAL) ($n=16/20$, 80%).

S. Typhimurium strains showed resistance to several antimicrobials, especially in relation to tetracycline ($n=11/11$), followed by ciprofloxacin ($n=10/11$), nalidixic acid, sulfonamides and chloramphenicol ($n=8/11$), streptomycin ($n=7/11$), trimethoprim-sulfamethoxazole ($n=6/11$), ampicillin ($n=4/11$), and gentamicin and cefotaxime ($n=1/11$) (Fig.1).

A total of *S. enterica* subsp. *enterica* 4,5,12: i: - strains ($n=7/7$) showed resistance to nalidixic acid, sulfamethoxazole trimethoprim-sulfonamides, tetracycline and streptomycin. For the other antimicrobials the resistant values of isolates were $n=6/7$ for ciprofloxacin, $n=5/7$ for chloramphenicol and $n=1/7$ to ampicillin, gentamicin and cefotaxime (Table 2).

Table 1. Number, frequency and serovars of *Salmonella* positive lymph nodes and feces from slaughtered pigs

Sample	N	Positives (%)	Serovars
Submandibular lymph nodes	50	10 (20%) ^a	- <i>S. enterica</i> subsp. <i>enterica</i> 4,5,12:i:- - <i>S. Typhimurium</i>
Mesenteric lymph nodes	50	9 (18%) ^a	- <i>S. Typhimurium</i> - <i>S. Derby</i> - <i>S. Brandenburg</i>
Feces	50	1 (2%) ^b	- <i>S. enterica</i> subsp. <i>enterica</i> 4,5,12:i:-
Mediastinal lymph nodes	50	0 (0%) ^b	-
TOTAL	200	20 (10%)	4 serovars

^{a,b} Different letters indicates statistical difference ($p < 0.05$) in the percentage of positives for each sample type.




Fig.1. Percentage of resistant *Salmonella* strains isolated from swine feces and lymph nodes for the tested antimicrobial compounds. Ciprofloxacin (CIP), Tetracycline (TET), Nalidixic Acid (NAL), Sulfonamides (SUL), Chloramphenicol (CLO), Streptomycin (EST), Sulfamethoxazole-trimethoprim (SUT), Ampicillin (AMP), Cefotaxime (CTX), Ceftriaxone (CRO), Gentamicin (GEN) Amikacin (AMI), Aztreonam (AZT), Cefepime (CPM), Cefoxitin(CFO), Ceftazidime (CAZ) Meropenem (MER).

Table 2. Antimicrobial resistance profile of different serotypes of *Salmonella* isolated from slaughtered pigs

Serovar	Antimicrobial
<i>S. Typhimurium</i>	CIP
<i>S. Brandenburg</i>	CIP, CRO
<i>S. Typhimurium</i>	NAL, CIP, CLO, TET
<i>S. Typhimurium</i>	CIP, SUT, SUL, TET
<i>S. Typhimurium</i>	NAL, CIP, CLO, EST, TET
<i>S. Typhimurium</i>	NAL, CTX, CIP, CLO, TET
<i>S. Typhimurium</i>	NAL, AMP, CIP, SUL, TET
<i>S. enterica</i> subsp. <i>enterica</i> 4,5,12:i:-	NAL, CIP, SUT, SUL, EST, TET
<i>S. Typhimurium</i>	AMP, CLO, SUT, SUL, EST, TET
<i>S. Typhimurium</i>	AMP, CIP, CLO, SUL, SUT, EST, TET
<i>S. Derby</i>	AMP, CIP, CLO, EST, SUL, SUT, TET
<i>S. Typhimurium</i>	NAL, CIP, CLO, SUT, SUL, EST, TET
<i>S. enterica</i> subsp. <i>enterica</i> 4,5,12:i:-	NAL, CIP, CLO, SUT, SUL, EST, TET
<i>S. enterica</i> subsp. <i>enterica</i> 4,5,12:i:-	NAL, CIP, CLO, SUT, SUL, EST, TET
<i>S. Typhimurium</i>	NAL, CIP, CLO, SUT, SUL, EST, TET
<i>S. enterica</i> subsp. <i>enterica</i> 4,5,12:i:-	NAL, CIP, CLO, SUT, SUL, EST, TET
<i>S. enterica</i> subsp. <i>enterica</i> 4,5,12:i:-	NAL, CIP, CLO, SUT, SUL, EST, TET
<i>S. Typhimurium</i>	NAL, CIP, CLO, SUT, SUL, EST, TET
<i>S. enterica</i> subsp. <i>enterica</i> 4,5,12:i:-	NAL, AMP, SUT, GEN, SUL, EST, TET
<i>S. enterica</i> subsp. <i>enterica</i> 4,5,12:i:-	NAL, CTX, CIP, CLO, SUT, SUL, EST, TET

Ciprofloxacin (CIP), Tetracycline (TET), Nalidixic acid (NAL), sulfonamides (SUL), chloramphenicol (CLO), streptomycin (EST), Trimethoprim-sulfamethoxazole (SUT), ampicillin (AMP), cefotaxime (CTX), ceftriaxone (CRO), Gentamicin (GEN).

The only one *S. Brandenburg* strain shows resistance exclusively to ciprofloxacin and ceftriaxone. The other single isolate of *S. Derby* was resistant against ampicillin, trimethoprim-sulfamethoxazole, sulfonamides, streptomycin, tetracycline, chloramphenicol and ciprofloxacin (Table 2).

Among the serovars identified, all were resistant to at least one of the antimicrobials tested, with 90% of them (n=18) resistant to at least four antimicrobial simultaneously. Considering the standards determined by CLSI for MDR samples, simultaneously resistant to ampicillin, chloramphenicol, streptomycin, trimethoprim-sulfamethoxazole and tetracycline, 15% (n=3) of the samples showed this pattern (Table 2). There was no strain positive for the ESBL enzyme production.

DISCUSSION

The present study revealed 10% frequency of *Salmonella* isolated from fecal and lymph nodes of pigs. This result is relatively lower to similar studies described worldwide. In Portugal, Vietnam and European Union were reported respectively 17.6%, 34.8% and 33% of positive identification of *Salmonella* in pigs (EFSA 2010, Ellerbroek et al. 2010, Gomes-Neves et al. 2012). Particularly in Brazil, different studies carried out in different regions revealed 16.6% and 67% of *Salmonella* from slaughtered pigs (Bessa et al. 2004, Sanchez et al. 2007, Silva et al. 2009, Kich et al. 2011).

Salmonella was most prevalent in the submandibular and mesenteric lymph nodes, then feces, and then mediastinal lymph nodes. It is known that the amount of feces sampled is related to the chances of the pathogen isolation (Davies et al. 1998), so the lower isolation rate in feces can be related to the sample aliquot used (1g). Other factor which can be associated to the lower frequency of *Salmo-*

nella in the fecal samples is the intermittent excretion of the pathogen. On the other hand, *Salmonella* is present intracellularly in the lymph nodes, which acts as reservoirs of the pathogen (Bahnon et al. 2006), being less exposed to other factors that would difficult the pathogen detection by the isolation technique, being the presence of *Salmonella* in the lymph nodes an indicator of the carrier status of the animal.

The differences observed in the isolation of *Salmonella* from specific lymph nodes relates to the anatomical position of them. The most prevalent occurrence of the pathogen was observed in the mesenteric and submandibular lymph nodes. The high prevalence of *Salmonella* in mesenteric and submandibular lymph nodes is associated with the proximity of them, respectively, to the gastrointestinal fecal contents, taking these nodes as an initial barrier to the pathogen. In many cases, the submandibular infection does not develop enteric signs, turning the animals into reservoirs. In fact, the oral-fecal cycle of *Salmonella* infections in pigs was investigated in other studies, showing that 70% of the isolates were identified in the tonsils and submandibular lymph nodes (Bahnon et al. 2006, Straw et al. 2012).

The high occurrence of *S. Typhimurium* identified in pigs sampled, agreed with increased prevalence of this serotype in the global scenery (Carrasco et al. 2012). In 2011, *S. Typhimurium* was also the predominant serovar reported in the United States, followed by *S. Enteritidis* (CDC 2013). The high prevalence of this serovar has been reported in European Union (EFSA 2010) and Brazil (Kich et al. 2011) as well.

Another relevant finding in the current study is 35% (n=7/20) of positive isolation of *S. enterica* subsp. *enterica* 4,5,12:i-, a serotype considered similar to *S. Typhimurium*, characterized by minor differences in the flagellar phase. Currently, *S. enterica* subsp. *enterica* 4,5,12:i- is referred as one of the main serotypes isolated from pigs worldwide, particularly in Europe (EFSA 2010) and the United States, since its notifications increased 351% between 2001 and 2011 (CDC 2013). This monophasic Typhimurium-like strains are considered an emerging pathogen, being associated to several outbreaks with high antibiotic resistance rates, however Brazilian data about this serovar isolation are still scarce.

The others serotypes also identified in the present study, as *S. Brandenburg* and *S. Derby*, only one sample each, have similar results to those observed in other countries (Kich et al. 2011, Carrasco et al. 2012, Thai et al. 2012).

In the current study, up 80% of strains were found to be resistant to ciprofloxacin, tetracycline and nalidixic acid. Similar studies in Brazil have also found a prevalence of resistance by *Salmonella* isolated from pigs to tetracycline (96.5%) and nalidixic acid (95.5%) (Castagna et al. 2001), as well as multidrug resistant strains particularly to sulfonamides (97.8%) and streptomycin (82.6%) (Weiss et al. 2002). The high resistance rates of our isolates for ciprofloxacin are similar to those found in pigs from Spain (97.1%) (EFSA 2010). In contrast, studies in other countries revealed minor occurrence of resistant *Salmonella* strains against tetracycline, sulfonamides and nalidixic

acid (Gomez-Laguna et al. 2011, Thai et al. 2012). Interestingly, ciprofloxacin, tetracycline and nalidixic acid that showed low effectiveness in our *Salmonella* strains, are considered antimicrobials of choice to humans and animal therapy of salmonellosis (Lesser & Miller 2005, Radostits et al. 2007).

The high prevalence antibiotic-resistant strains of *Salmonella* encountered in our study is similar to the reported in other studies, such as investigations in pigs from Vietnam, Estonia, Ireland and Brazil. These studies reported multidrug resistance frequencies of 23.2%, 13.6%, 59.4% and 24.2%, respectively (EFSA 2010, Thai et al. 2012)

Multidrug resistant bacteria, including *Salmonella* species from animal and human origin is an emergent public health concern (Giguère et al. 2010). The high antimicrobial resistance rates in pig production can be attributed, among other reasons, to the improper use of antimicrobials for treatment of diseases or growth promotion in pork production (Gebreyes et al. 2004, Kich et al. 2011, Gomes-Neves et al. 2012). However, some resistance mechanisms to some antimicrobials are easily transferred between species, and their prevalence may be in part be due to environmental contamination. Another important resistance acquisition route are the genes that promotes simultaneous resistance to different compounds. In the light of this, ubiquitous and opportunistic pathogenic bacteria as *Salmonella* with resistance to conventional antimicrobials can survive in environment of farms, and are able to transmit drug resistance to other bacteria that infect wildlife and domestic animals, including pigs (Thakur et al. 2007). Indeed, despite the prohibition of the use of chloramphenicol in livestock 25 years ago in United States (Thakur et al. 2007) and 18 years ago in Brazil (Brasil 2003), a diverse range of bacteria remain resistant to these drugs even today.

In our study were not detected *Salmonella* strains ESBL-enzyme positive. However, in Europe ESBL production was referred in 0.6% of the strains isolated from pigs, when tested for cefotaxime and 0.5% for ceftazidime (EFSA 2014). Despite absence of *Salmonella* strains positive for ESBL-production in pigs sampled, the emergence of this pathogenic mechanism of antimicrobial resistance of *Salmonella*, requires continuous epidemiological vigilance studies.

In this study, we found that *Salmonella* spp. isolated from pigs exhibit high levels of resistance to antimicrobials. These antibiotic-resistant strains may potentially cause foodborne outbreaks of antibiotic-resistant salmonella in humans. *S. Typhimurium* was the most common serovar, even with the adoption of control measures implemented over several decades in pig farms. In addition, there is an increasing occurrence of *S. enterica* subsp. *enterica* 4,5,12:i:-, considered an emergent human pathogen and with few Brazilian data about this serovar available. The high resistance rates found in our study reveals that despite regional differences for resistance to some antimicrobials, the inappropriate use of antimicrobials increases the selection rate of multidrug resistant bacteria, including *Salmonella*. Thus, the selection of first-choice antimicrobial treatment should be based on local *in vitro* resistance patterns. Indeed, the responsible use of antimicrobials for animals is an

emergent One Health concern, to conserve these drugs for human therapy approaches.

Acknowledgements.- To Adolfo Lutz Institute for conducting the serotyping of strains and Cefar Diagnostic for providing the antibiogram discs.

Conflict of Interest.- No conflict of interest.

REFERENCES

- ABIPECS 2013. Produção mundial de carne suína. Associação Brasileira da Indústria Produtora e Exportadora de Carne Suína.
- Andrews W.H., Jacobson A. & Hammack 2014. BAM Chapter 5: *Salmonella*, p.5.01-05.020. In: Food and Drug Administration (Ed.), Bacteriological Analytical Manual. 8th ed. AOAC International, Gaithersburg.
- Bahnon P.B., Damman D.J., Isaacson R.E., Miller G.Y. & Weigel R.M. 2006. Prevalence and serovars of *Salmonella enterica* isolated from ileocolic lymph nodes of market pigs reared in selected Midwest US swine herds. J. Swine Health Prod. 4:182-188.
- Bessa M.C., Costa M.D. & Cardoso M. 2004. Prevalência de *Salmonella* sp. em suínos abatidos em frigoríficos do Rio Grande do Sul. Pesq. Vet. Bras. 24:80-84.
- Bollaerts K., Aerts M., Faes C., Grijspeerd K., Dewulf J. & Mintiens K. 2008. Human salmonellosis: estimation of dose-illness from outbreak data. Risk Analysis 28:427-440.
- Brasil 2003. Instrução Normativa nº 9, de 27 de junho de 2003. Ministério da Agricultura Pecuária e Abastecimento (MAPA).
- Carrasco E., Morales-Rueda A. & Garcia-Gimeno R.M. 2012. Cross-contamination and recontamination by *Salmonella* in foods: a review. Food Res. Int. 45:545-556.
- Castagna S.M.F., Bessa M.C., Carvalho D.A., Cardoso M. & Costa M. 2001. Resistência a antimicrobianos de amostras de *Salmonella* sp. isoladas de suínos abatidos no estado do Rio Grande do Sul. Arq. Fac. Vet. UFRGS 29:44-49.
- CDC 2013. Centers for Disease Control, Prevention National *Salmonella* Surveillance Annual Report - Appendices, 2011. US Department of Health and Human Services, CDC, Atlanta, GA.
- CDC 2014. Centers for Disease Control, Human Isolates Final Report, 2012, National Antimicrobial Resistance Monitoring System for Enteric Bacteria (NARMS), p.12-26.
- CSLI 2013. Performance standards for antimicrobial susceptibility testing M100-S23. Clinical Laboratory Standards Institute, Wayne, PA.
- Davies P.R., Bovee G.E.M., Funk J.A., Morrow W.E.M., Jones F.T. & Deen J. 1998. Isolation of *Salmonella* serotypes from feces of pigs raised in a multiple-site production system. J. Am. Vet. Med. Assoc. 212:1925-1929.
- Duggan S.J., Mannion C., Prendergast D.M., Leonard N., Fanning S., Gonzales-Barron U., Egan J., Butler F. & Duffy G. 2010. Tracking the *Salmonella* status of pigs and pork from Lairage through the slaughter process in the Republic of Ireland. J. Food Prot. 73:2148-2160.
- EFSA 2010. The community summary report on trends and sources of zoonoses, zoonotic agents and food-borne outbreaks in the European Union in 2008. European Food Safety Authority, EFSA J. 8(1):23-110.
- EFSA 2014. The European Union Summary Report on antimicrobial resistance in zoonotic and indicator bacteria from humans, animals and food in 2012. European Food Safety Authority, EFSA J. 12:23-51.
- Ellerbroek L., Narapati D., Phu Tai N., Poosaran N., Pinthong R., Sirimalaisuan A., Tshering P., Fries R., Zessin K.H., Baumann M. & Schroeter A. 2010. Antibiotic resistance in *Salmonella* isolates from imported chicken carcasses in Bhutan and from pig carcasses in Vietnam. J. Food Prot. 73:376-379.
- Gebreyes W.A., Thakur S., Davies P.R., Funk J.A. & Altier C. 2004. Trends in antimicrobial resistance, phage types and integrons among *Salmonella* serotypes from pigs, 1997-2000. J. Antimicrob. Chemother. 53:997-1003.
- Giguère S., Prescott J.F., Baggot J.D., Walker R.D. & Dowling P.M. 2010. Terapia antimicrobiana em medicina veterinária. Roca, São Paulo. 683p.

- Gomes-Neves E., Antunes P., Tavares A., Themudo P., Cardoso M.F., Gartner F., Costa J.M. & Peixe L. 2012. *Salmonella* cross-contamination in swine abattoirs in Portugal: carcasses, meat and meat handlers. *Int. J. Food Microbiol.* 157:82-87.
- Gomez-Laguna J., Hernandez M., Creus E., Echeita A., Otal J., Herrera-Leon S. & Astorga R.J. 2011. Prevalence and antimicrobial susceptibility of *Salmonella* infections in free-range pigs. *Vet. J.* 190:176-178.
- Grimont P.A.D. & Weil F.X. 2007. Antigenic Formulae of the *Salmonella* Serovars. Institut Pasteur, Centre Collaborater OMS de Référence et de Recherche pour les *Salmonella*, Paris, France. 166p.
- Kich J.D., Coldebella A., Mores N., Nogueira M.G., Cardoso M., Fratamico P.M., Call J.E., Fedorka-Cray P. & Luchansky J.B. 2011. Prevalence, distribution, and molecular characterization of *Salmonella* recovered from swine finishing herds and a slaughter facility in Santa Catarina, Brazil. *Int. J. Food Microbiol.* 151:307-313.
- Lesser F.C. & Miller I.S. 2005. Diseases caused by Gram-negative bacteria, p.901. In: Harrison T.K.D.L., Fauci A.S., Longo D.L., Braunwald E., Hauser S.L. & Jameson J.L. (Eds), *Principles of Internal Medicine*. McGraw-Hill Education, New York.
- Letellier A., Beauchamp G., Guevremont E., D'Allaire S., Hurnik D. & Quessy S. 2009. Risk factors at slaughter associated with presence of *Salmonella* on hog carcasses in Canada. *J. Food Prot.* 72:2326-2331.
- Radostits O.M.G., Gay C.C., Hinchcliff K.W. & Constable P.D. 2007. *Veterinary Medicine: A textbook of the diseases of cattle, horses, sheep, pigs, and goats*. Saunders Elsevier, Philadelphia. 2156p.
- Sanchez J., Dohoo I.R., Christensen J. & Rajic A. 2007. Factors influencing the prevalence of *Salmonella* spp. in swine farms: a meta-analysis approach. *Prev. Vet. Med.* 81:148-177.
- Santos I.R., Nascimento V.P. & Flores M.L. 2002. *Salmonella enteritidis* isoladas de amostras clínicas de humanos e de alimentos envolvidos em episódios de toxi-infecções alimentares, ocorridas entre 1995 e 1996, no Estado do Rio Grande do Sul. *Hig. Aliment.* 16:93-99.
- Silva M.C., Faria G.S., Paula D.A.J., Martins R.P., Caramori Junior J.G., Kich J.D., Colodeli E.M., Nakazato I. & Dutra V. 2009. Prevalência de *Salmonella* sp. em suínos abatidos no Estado de Mato Grosso. *Ciência Rural* 39:266-268.
- Sousa Junior M.A., Ferreira E.S. & Conceição G.C. 2004. Betalactamases de espectro ampliado (ESBL): um importante mecanismo de resistência bacteriana e sua detecção no laboratório clínico. *NewsLab.* 63:153-174.
- Straw B.E., Zimmerman J.J., Karriker I.A., Ramirez A., Schwartz K.J. & Stevenson G.W. 2012. *Diseases of Swine*. 10th ed. Wiley-Blackwell, London.
- Thai T.H., Hirai T., Lan N.T. & Yamaguchi R. 2012. Antibiotic resistance profiles of *Salmonella* serovars isolated from retail pork and chicken meat in North Vietnam. *Int. J. Food Microbiol.* 156:147-151.
- Thakur S., Tadesse D.A., Morrow M. & Gebreyes W.A. 2007. Occurrence of multidrug resistant *Salmonella* in antimicrobial-free (ABF) swine production systems. *Vet. Microbiol.* 125:362-367.
- Weiss L.H., Nonig R.B., Cardoso M. & Costa M.D. 2002. Ocorrência de *Salmonella* spp. em suínos de terminação no Rio Grande do Sul. *Pesq. Vet. Bras.* 22:104-108.