

ANAIS



Simpósio Internacional de Nutrição e Saúde de Peixes

25 a 27 de outubro de 2017

UNESP - FAZENDA EXPERIMENTAL
LAGEADO BOTUCATU - SP

Apoio

Realização



AquaNutri
FMVZ - UNESP - BOTUCATU

FICHA CATALOGRÁFICA ELABORADA PELA SEÇÃO TÉC. AQUIS. TRATAMENTO DA INFORM.
DIVISÃO DE BIBLIOTECA E DOCUMENTAÇÃO - CAMPUS DE BOTUCATU - UNESP
BIBLIOTECÁRIA RESPONSÁVEL: **ROSEMEIRE APARECIDA VICENTE**

Simpósio Internacional de Nutrição e Saúde de Peixes (7 : 2017 : Botucatu)
Anais [do] VII Simpósio Internacional de Nutrição e Saúde de Peixes, 25 a 27
de outubro de 2017 / Organizadores Luiz Edivaldo Pezzato, Margarida Maria
Barros, José Eurico Possebon Cyrino, Wilson Massamitu Furuya. - Botucatu :
UNESP-FCA, 2017

Resumos

1. Peixe - Nutrição. 2. Peixe – Saúde. 3. Fenômenos fisiológicos da nutrição. 4.
Nutrição animal. 5. Resumos. 6. Universidade Estadual Paulista "Júlio de
Mesquita Filho", Fazenda Experimental Lageado. I. Título. II. Pezzato, Luiz
Edivaldo. II. Barros, Margarida Maria. III. Cyrino, José Eurico Possebon.
IV. Furuya, Wilson Massamitu.

CDD 639.3

Resumos apresentados durante o VII Simpósio Internacional de Nutrição e Saúde de Peixes



Nutritional quality of shrimp *Litopenaeus vannamei* byproducts meal

Sidely Gil Alves Vieira¹, Fabíola Helena dos Santos Fogaça^{*}, Luiz Gonzaga Alves dos Santos Filho¹, Alitiane Moura Lemos Pereira², João Avelar Magalhães², Irani Alves Ferreira³, Thiago Fernandes Alves Silva⁴ *Pesquisador científico; Embrapa Meio-Norte; Br 343 km 35, 64200-970 – Parnaíba – PI; fabiola.fogaca@embrapa.br; ¹Universidade Federal do Piauí, Parnaíba, PI; ² Embrapa Meio-Norte, Parnaíba, PI; ³University of Maryland, College Park, MD, USA; ⁴CAUNESP, Jaboticabal, SP

The use of alternative food to those traditionally used in commercial diets is a growing demand from the aquaculture sector. At the same time, the processing industries seek technologies to use waste from seafood processing in order to minimize their environmental impact and add value to byproducts. The shrimp *Litopenaeus vannamei* is the principal seafood produced by Brazilian Northeast aquaculture and its processing generates lot of byproducts (50% of total biomass) that could be used as diet ingredient for aquatic organisms. The aim of this study was to prepare a shrimp byproducts meal and evaluate its nutritional quality. Shrimp byproducts (120 kg) were obtained from a seafood processing industry of Ceará, Brazil. The marine shrimp head and carapace were washed with chlorinated water, drained, homogenized in electric grinder and dried in a forced air circulation oven at 60 °C for 24 hours. After drying, the material was weighed, minced with knife mill, sieve hole of 1.5 mm diameter, packed in polyethylene bags and stored in a freezer (- 20 ° C). The nutritional quality was determined by analysis of crude protein (CP), essential amino acids, ether extract (EE), fatty acids, mineral matter (MM), Calcium (Ca), Phosphorus (P), Iron (Fe), Copper (Cu) and Zinc (Zn). The meal showed high protein content with 55.18% CP. The protein quality was evaluated by its essential amino acids profile: 3.43 % for lysine, 1.05 % for methionine, 2.05 % for methionine+cysteine, 1.74 % for threonine, 2.80 % for arginine, 2.42 % for phenylalanine, 1.09 % for histidine, 1.97 % for isoleucine, 3.09 % for leucine, 0.50 % for tryptophan and 2.54 % for valine. EE levels summarizing 6.25 %, of saturated fatty acids and 1.56 % of polyunsaturated fatty acids (PUFA). Linoleic acid (ω -6) showed a content of 0.80 % and arachidonic acid showed 0.08 % content. There was no detection of linolenic (ω -3), eicosapentaenoic (EPA) and docosahexaenoic (DHA) acids. The MM showed levels of 19.00 %, 5.44 % of Ca and 1.23 % of P, higher than recommended for fish feed (15.00 % MM and 0.75 % P). Cu (70.08 mg.kg⁻¹) and Fe (80.79 mg.kg⁻¹) attend more than 100 % of the fish requirements, while Zn (71.97 mg.kg⁻¹) provides 90.52 %. According to the results, the shrimp byproducts meal was not a good source of long chain fatty acids (EPA, DHA and arachidonic), which are ω -3 and ω -6 precursors, but showed a good nutritional quality protein, which provides good levels of essential amino acids. However, it has a high mineral content, a factor which may limit its inclusion in commercial diets. Future researches should be to carry out to assess the levels of shrimp byproducts meal inclusion in diets for fish.