

## Root Distribution of Four Tropical Useful Plants after Six Months of Cultivation on Degraded Sites in Amazonia

K. Voß<sup>1</sup>, L. Gasparotto<sup>2</sup>, R. Lieberei<sup>1</sup> and F. Feldmann<sup>1</sup>

<sup>1</sup>Institut für Angewandte Botanik der Universität Hamburg

<sup>2</sup>EMBRAPA/CPAA, C.P. 319, km 28, AM 010, Manaus-AM, Brazil

The knowledge about the genetically and environmentally influenced root formation pattern of tropical useful plants is of great importance for adapting field management processes. Especially in the installation phase of a plantation of perennial plants the characteristics of each plant species concerning root growth, spread and distribution, its strategy of nutrient uptake as well as its velocity of mycorrhiza formation can decide on the ability of the plant to establish itself at degraded sites. Moreover these qualities are of interest for the species' usefulness to cover previously burned and cleared soils, to avoid erosion and minimise nutrient losses.

Perennial plants commonly are precultivated in nurseries for six to eight weeks. The environmental factors of nursery pots or plastic bags are characterised by low volume, adequate substrate, and equilibrated regimes of fertilisation and irrigation. Additionally the plants in the pots can be inoculated with high contents of effective arbuscular mycorrhiza fungi (AMF) of commonly exogenic genotypes.

In the field the plants are set into holes, fertilised in graduated levels, and then surrounded by organic material and disturbed soil of the B-horizon, both containing few infection units of native AMF. There, in the early stages of development the plant material has to overcome drastic environmentally changes as soon as the roots are growing into the adjacent soil. This relatively undisturbed field soil is characterised by chemical and physical degradation, e. g. low nutrient availability, low pH and often compactness. Here the roots meet the first time larger quantities of native AMF which potentially will form now the larger part of the symbioses of the roots with AMF.

How do such changes of the environment affect the root distribution of crop species? In order to put forward that question we selected four species which are presumed to form different types of root systems for preliminary investigations of the root spread: *Bixa orellana* L., *Bixaceae*, *Bertholletia excelsa* HUMB. ET BONPL., *Lecythidaceae*, *Theobroma grandiflorum* (WILLD. EX SPRENG) SCHUM., *Sterculiaceae* and *Hevea brasiliensis* MUELL. ARG., *Euphorbiaceae*. Before planting and six months after planting in the experimental field near Manaus, Amazonia, the root system and root distribution throughout the soil horizons were described. The nutrient status of the field soil was analysed. The fine roots were characterised by morphological parameters to enable detection and classifying under field conditions. Fertiliser induced differences in root distribution were described for *Bixa orellana*.

The results are discussed with regard to the species' rooting strategies, the spatial and temporal coincidence of the mycorrhizal symbionts in the field and the common management practice. An outlook is made for necessary further quantitative and qualitative examinations to prove the preliminary conclusions.