

EFFECT OF AGE OF A SORGHUM-SUDANGRASS HYBRID ON ITS ALLELOPATHIC ACTION

Efeito da Idade de um Híbrido de Sorgo com Capim-Sudão em sua Ação Alelopática

MARCHI, G.², MARCHI, E.C.S.³, WANG, G.⁴ and MCGIFFEN, M.⁵

ABSTRACT - Hybrids of *Sorghum sudanensis* (sudangrass) and *Sorghum bicolor* genotypes can produce high amounts of biomass, sorgoleone (a long chain hydroquinone), and other phytotoxic substances. Shoots and roots of a sorghum-sudangrass hybrid (cv. Trudan 8) were collected 10, 20, 30, 40, and 50 days after emergence. Four concentrations of aqueous extracts from the shoots and roots (0, 0.4, 2, and 10 g L⁻¹, w/v) were used to treat seeds of lettuce (*Lactuca sativa*), tomato (*Lycopersicum sculentum*), purslane (*Portulaca oleracea*), and pigweed (*Amaranthus retroflexus*). Seed germination of lettuce, tomato, and pigweed was inhibited by extracts from sorghum-sudangrass shoots at 10 g L⁻¹ when made from sorghum-sudangrass plants 20 days or less in age. Seed germination of purslane was not inhibited by any sorghum-sudangrass extract. Growth of the four species evaluated were systematically inhibited when treated with 10 g L⁻¹ extracts from sorghum-sudangrass shoots harvested up to 10 days after emergence.

Keywords: allelopathy, germination inhibition, lettuce (*Lactuca sativa*), tomato (*Lycopersicum sculentum*), pigweed (*Amaranthus retroflexus*) and purslane (*Portulaca oleracea*).

RESUMO - Os capins híbridos obtidos pelo cruzamento entre *Sorghum sudanensis* (capim-sudão) e genótipos de *Sorghum bicolor* possuem alto potencial para produção de biomassa e para controle de plantas daninhas pela produção de substâncias fitotóxicas, como o sorgoleone (uma hidroquinona de cadeia longa). Sementes de alface (*Lactuca sativa*), tomate (*Lycopersicum sculentum*), beldroega (*Portulaca oleracea*) e caruru (*Amaranthus retroflexus*) foram submetidas a tratamentos com extratos aquosos da parte aérea e das raízes do híbrido de sorgo com capim-sudão, cv. Trudan 8, colhido em cinco diferentes estádios de crescimento (10, 20, 30, 40 e 50 dias após a emergência). Os extratos foram preparados em quatro concentrações (0, 0,4, 2 e 10 g L⁻¹, p/v) e aplicados em quatro repetições. Após os tratamentos, a germinação e o comprimento de plântulas das espécies foram avaliados. A germinação de sementes de tomate, caruru e alface foi inibida pelos extratos da parte aérea das plantas de Trudan 8, na concentração de 10 g L⁻¹, colhidas até os 20 dias após a emergência. A germinação de sementes de beldroega, no tocante à porcentagem de germinação, não foi inibida pelos extratos de Trudan 8. O crescimento das quatro espécies avaliadas foi inibido quando tratadas com extratos aquosos da parte aérea de Trudan 8, colhida até os 10 dias após a emergência, na concentração de 10 g L⁻¹.

Palavras-chave: alelopatia, inibidor de germinação, alface (*Lactuca sativa*), tomate (*Lycopersicum sculentum*), caruru (*Amaranthus retroflexus*) e beldroega (*Portulaca oleracea*).

¹ Recebido para publicação em 19.5.2008 e na forma revisada em 19.9.2008.

² Pesquisador, Embrapa Cerrados; Rodovia BR020, km 18, Caixa Postal 08223, 73310-970, Planaltina-DF, <giuliano.marchi@cpac.embrapa.br>; ³ Bolsista, CNPq/Embrapa Cerrados; ⁴ Pesquisador, Maricopa Agricultural Center, University of Arizona, 37860 W. Smith-Enke Road, Maricopa, AZ 85239, USA; ⁵ Prof., Department of Botany and Plant Sciences, University of California, Riverside, CA 92521, USA



INTRODUCTION

Both organic and conventional farmers apply no tillage systems. Among the crops used for pasture and straw mulch for integrated crop-livestock production, cutting or foraging sorghum-sudangrass hybrids produce large amounts of biomass, suppress weeds, and decrease soil compaction (McKinney et al., 2004). Sorghum-sudangrass cultivars are produced from crossing inter-specific *Sorghum sudanensis* Piper Stapf hybrids with *Sorghum bicolor* genotypes (Raupp & Brancão, 2000). Growing these hybrids without irrigation during February in Central Brazil has yielded 9 to 12 tons of dry matter in three successive cuts, for a total of at least 40 to 60 tons of fresh forage per hectare (Zago, 1997; Mello et al., 2003). Sorghum-sudangrass hybrids also present high forage production potential when grown at the beginning of the rainy season in Central Brazil (Tomich et al., 2004).

Besides its high biomass production potential, sorghum-sudangrass hybrids release phytotoxic substances into the environment, being able to suppress weed development in some situations (Forney & Foy, 1985). Sorghum species produce a myriad of substances with allelopathic potential that may inhibit weed germination and growth (Weston, 1996). The allelopathic compounds are a secondary metabolism product, and their chemical compositions and concentrations vary according to species, age, and climatic conditions (Dayan, 2006). Using allelopathic substance-producing plants may allow farmers to reduce herbicide use (Weston, 1996; Bärberi & Mazzoncini, 2001).

Allelopathic effects of sorghum-sudangrass may be related to phenolic compounds released by tissue decomposition (Weston et al., 1989; Sene et al., 2001). Phenolic acids such as ferulic, *p*-coumaric, syringic, vanilic, and *p*-hydroxybenzoic are produced from sorghum hybrid roots during residue decomposition. These substances are responsible for growth inhibition of several plant species (Roth et al., 2000). Another compound found in sorghum genotypes is dhurrin, a cyanogenic glycoside that degrades to *p* hydroxybenzaldehyde, HCN, and glucose (Nicollier et al., 1983).

Netzley & Butler (1986) verified that sorghum roots exude the long chain

hydroquinone sorgoleone, which exhibits high phytotoxicity. This substance is a potent electron transport inhibitor in photosystem II in chloroplasts (González et al., 1997; Hejl & Koster, 2004). Sudangrass also produces from 1.3 to 1.9 mg sorgoleone g⁻¹, an amount similar to that produced by sorghum (Bais et al., 2006). Sorgoleone is active at very small concentrations (McGuire, 2003) and has a higher specific activity than diuron, one of the most potent photosynthetic inhibitors (Gonzalez et al., 1997).

Sorgoleone is rapidly degraded in soil (Czarnota et al., 2001), suggesting that large amounts must be continuously produced to maintain phytotoxicity in the soil solution (Bais et al., 2006). However, the capacity to produce plant allelochemicals declines as the plant ages (Woodhead, 1981; Weston et al., 1989; Ben-Hammouda et al., 1995; An et al., 2003), thus sorgoleone is mainly produced in younger plants (Dayan, 2006).

The aim of this work was to evaluate inhibition of lettuce, tomato, purslane, and pigweed seed germination and growth after treatment with shoot and root aqueous extracts of the sorghum-sudangrass hybrid cv. Trudan 8 harvested on five dates after germination.

MATERIALS AND METHODS

Extraction procedure

Seeds of sorghum (*Sorghum bicolor*) and sudangrass (*Sorghum sudanensis*) hybrid, cv. Trudan 8 (hereafter Sorghum-sudangrass), were planted in 32 pots under glasshouse conditions at the University of California, Riverside from August to November 2003. Average daily temperature during the experiment varied from 13 to 28 °C.

Plants were harvested at five ages: 10, 20, 30, 40, and 50 days after emergence. After each harvest, plants were washed with distilled water, separated as shoots and roots, and dried in oven at 60 °C. Shoots and roots were ground to a fine powder and extracted with distilled water for 24 h at 24 °C in a lit room. These extracts were filtered through four tissue sheets to remove particles and

filtered again with Whatman n.42. Four concentrations (0, 0.4, 2, and 10 g L⁻¹ w/v) of each extract were prepared and stored in the dark for 7 days at 4°C for further analysis. Electrical conductivity of the extracts ranged from 35 µS/cm to 1.51 mS/cm in the shoot extracts and from 15 µS/cm to 1.32 mS/cm in the root extracts, while pH ranged from 5.64 to 6.59 in the shoots and from 4.11 to 6.77 in the roots. These values were within the normal range and did not harm the germination or development of the plants (Everitt et al., 1983).

Seed germination and plant growth

Germination of lettuce (*Lactuca sativa*) cv. "Black Seed Simpson", tomato (*Lycopersicon esculentum*) cv. "Advantage 81376", purslane (*Portulaca oleracea*), and pigweed (*Amaranthus retroflexus*) seeds was measured after treatment with sorghum-sudangrass root and shoot extracts. The experiment was arranged in a completely randomized design with four replicates. For each replicate, thirty seeds of each species were placed into two petri dishes lined with sterile filter paper (Whatman #1). Seven mL of distilled water (control) or one of the four sorghum-sudangrass extracts were added to each Petri dish, totaling five experimental treatments. Petri dishes were sealed to prevent humidity loss or contamination, and kept in a lit room for 10 h followed by 14 h in the dark at 24 °C. Lettuce, tomato, purslane, and pigweed seed germination and plant growth were measured 4,5, 6, and 7 days after sowing, respectively. Seeds were counted as germinated after radicle emergence of at least 1 mm. Seven plants were randomly chosen from each Petri dish and their size measured.

Data analysis

Germination data were submitted to analysis at 5% probability confidence intervals. The data were binomially distributed and analyzed by calculating exact confidence intervals (Pratt, 1968; Blyth, 1996; Leemis & Trivedi, 1996) using Sisvar software (Ferreira, 2000). The statistical model used was $Y/n \sim \text{Binomial}("n, Y)$, where: Y denotes the number of successes (germinated seeds), n denotes

the number of observations (total seeds), and "n", the proportion of population successes for each sample.

Plant growth data were subjected to analysis of variance and the means were separated by the Tukey's test (5% probability).

RESULTS AND DISCUSSION

Seed germination

Seed germination of lettuce, tomato, and pigweed was inhibited by extracts from Sorghum-sudangrass plants harvested at 10 and 20 days after emergence, at 10 g L⁻¹ (Table 1). Root and shoot extracts prepared with Sorghum-sudangrass plants harvested 30, 40, and 50 days after emergence did not inhibit germination of these three species, suggesting that the allelopathic potential of sorghum-sudangrass plants harvested 30 days after germination decreased.

Lettuce seed germination was inhibited when using 2 and 10 g L⁻¹ sorghum-sudangrass shoot and root extracts from plants harvested 10 days after emergence. Extracts of sorghum-sudangrass prepared with plants harvested 20 to 50 days after emergence did not inhibit lettuce seed germination, except 10 g L⁻¹ extracts of sorghum-sudangrass shoot harvested 20 days after emergence, which did inhibit lettuce seed germination. Tomato and pigweed seed germination percentage was strongly reduced following application of 10 g L⁻¹ extracts of sorghum-sudangrass shoot harvested 10 or 20 days after emergence. Purslane seed germination was not inhibited by extracts of sorghum-sudangrass shoot or root.

Plant growth

As sorgoleone produced by sorghum-sudangrass affects photosystem II (González et al., 1997; Hejl et al., 2004), it is expected that the growth of other plants be affected intensely just after they germinate and start to photosynthesize (Barbosa et al., 1998). The effects of sorghum-sudangrass extracts on plant growth depended on the type of extract: root or shoot, its concentration, and its age at



Table 1 - Lettuce, tomato, purslane, and pigweed germination (%) after treatment with root or shoot aqueous extracts of sorghum-sudangrass harvested at 5 growth stages[§]

Extract concentration (g L ⁻¹)	Sorghum-sudangrass age (days after emergence)				
	10	20	30	40	50
	Germination (%)				
	Lettuce				
0	95.00 Aa	95.00 Aa	94.58 Aa	95.83 Aa	94.17 Aa
0.4R*	95.42 Aa	94.17 Aa	87.08 Aab	95.83 Aa	95.83 Aa
2R	83.33 ABab	94.58 Aa	87.08 Aa	93.33 Aa	92.50 Aa
10R	82.92 ABab	92.08 Aa	83.75 ABab	95.00 Aa	95.83 Aa
0.4S	91.25 Aa	94.17 Aa	87.08 Aab	95.83 Aa	96.67 Aa
2S	82.92 ABab	95.00 Aa	88.75 Aa	93.33 Aa	95.83 Aa
10S	64.58 Cb	84.58 ABa	89.58 Aa	92.50 Aa	91.67 Aa
	Tomato				
0	66.67 Aa	76.25 Aa	72.50 Aa	70.00 Aa	66.67 Aa
0.4R	70.41 Aa	69.17 Aa	68.75 Aa	75.83 Aa	67.50 Aa
2R	75.00 Aa	71.67 Aa	66.67 Aa	73.33 Aa	72.50 Aa
10R	70.00 Aa	68.10 Aa	71.66 Aa	68.33 Aa	71.67 Aa
0.4S	65.00 Aa	71.25 Aa	65.42 Aa	75.83 Aa	67.50 Aa
2S	60.83 Aa	73.75 Aa	68.33 Aa	69.17 Aa	68.33 Aa
10S	12.08 Bb	3.75 Bc	65.00 Aa	71.67 Aa	66.67 Aa
	Purslane				
0	81.25 Aa	75.00 Aa	78.75 Aa	72.50 Aa	78.33 Aa
0.4R	79.17 Aa	80.83 Aa	78.33 ABa	82.50 Aa	76.67 Aa
2R	81.67 Aa	81.67 Aa	80.83 Aa	69.17 ABb	82.50 Aa
10R	79.17 Aa	80.83 Aa	78.33 ABa	80.00 Aa	77.50 Aa
0.4S	79.17 Aa	74.17 Aa	80.83 Aa	75.83 Aa	76.67 Aa
2S	80.00 Aa	74.58 Aab	89.17 Aa	78.33 Aab	80.83 Aa
10S	76.25 Aa	75.42 Aa	74.17 ABa	73.33 Aa	84.17 Aa
	Pigweed				
0	90.00 Aa	91.67 Aa	92.08 Aa	90.00 ABa	96.67 Aa
0.4R	89.58 Aa	88.33 Aa	90.42 Aa	91.67 ABa	89.17 ABa
2R	87.50 Aa	86.25 Aa	83.33 Aab	94.17 Aa	88.33 ABa
10R	89.05 Ab	87.50 Ab	88.33 Ab	98.33 Aa	90.00 Aa
0.4S	80.42 Aab	86.67 Aab	89.58 Aa	96.67 Aa	93.33 Aa
2S	82.08 Aab	85.42 Aab	89.58 Aab	97.50 Aa	95.83 Aa
10S	2.92 Bb	2.08 Bb	85.83 Aa	95.00 Aa	90.00 Aa

§ Different concentrations (0.4, 2, 10 g L⁻¹) of root (R) or shoot (S) extracts from sorghum-sudangrass plants harvested at 5 different stages (10, 20, 30, 40, 50 days after germination) were used to treat the seeds of four plant species (lettuce, tomato, purslane, and pigweed). For each species, plant length in the same column followed by the same upper case letter was not significantly different, while values in the same rows followed by the same lower case letter were not significantly different by the Tukey test $p \leq 5\%$.

* R = root extracts; S = shoot extracts.

harvest (Table 2). Shoot extracts inhibited growth more strongly than those prepared from roots. This result is consistent with Correia et al. (2005a), who showed that leaf extracts from two sorghum genotypes, XBG00478 and DKB860, inhibited soybean radicle length more than the root extracts.

Extracts of 10 g L⁻¹ sorghum-sudangrass prepared from shoot harvested at 10 and 20 days after emergence strongly inhibited the growth of all the plant species (Table 2). Sorghum-sudangrass 2 and 10 g L⁻¹ extracts prepared from shoot or root harvested 10 days after emergence, inhibited lettuce, tomato, purslane, and pigweed plant length. However, 10 g L⁻¹ extracts prepared with shoots strongly inhibited seedling growth, mainly when shoot was harvested 10 and 20 days after emergence. Tomato and pigweed seedlings either grew poorly or did not grow at all when treated with 10 g L⁻¹ extracts prepared from shoots harvested 10 and 20 days after emergence.

Correia et al. (2005a) suggest that radicle inhibition by allelopathic substances in Petri dish experiments is more likely to occur than shoot inhibition, as absorption and concentration of allelochemicals by root tissues are favored by physical contact with the filter paper. In a field experiment, Xuan et al. (2004) determined that plant leaves produce more allelochemicals and thus have higher allelopathic potential than roots or stems. We found that both shoots and roots were strongly inhibited by sorghum-sudangrass extracts (Tables 3 and 4).

Lettuce, tomato, purslane, and pigweed radicle and shoot growth was strongly inhibited following treatment with 0.4, 2 and 10 g L⁻¹ sorghum-sudangrass shoot extracts, especially when the extracts were prepared with sorghum-sudangrass harvested 10 days after emergence (Table 3). Any of the 10 g L⁻¹ sorghum-sudangrass shoot extracts other than those prepared from 30-day-old plants inhibited radicle growth of all the target species. Tomato and pigweed roots and shoots either grew minimally or not at all when treated with

10 g L⁻¹ extracts prepared with shoots harvested 10 and 20 days after emergence.

Sorghum-sudangrass extracts inhibited both seed germination and growth. Ten g L⁻¹ sorghum-sudangrass extracts inhibited lettuce, tomato, purslane and pigweed germination and root and shoot growth, mainly when sorghum-sudangrass shoot extracts were prepared from 10-day-old plants. This result is important mainly when sorghum-sudangrass is planted with other crops in crop-livestock integration and affects the productions of other crops (Correia et al., 2005b; Olibone et al., 2006).

Since older sorghum-sudangrass tissue presents a lower concentration of allelopathic chemicals, high biomass production does not always mean high allelopathic potential. These results suggest that sorghum-sudangrass allelochemicals are produced at a higher intensity up to 10 days after its emergence, and are found primarily in the shoots. The allelopathic potential of aqueous extracts from shoots of the hybrid Sudex [(*Sorghum bicolor*) Moench x *Sorghum Sudanese* (Piper) Stapf], cv. FFR 201] peaks 7 days after germination (Weston et al., 1989). In the present experiment, extracts prepared with sorghum-sudangrass harvested at 30, 40, and 50 days after emergence did not inhibit seed germination (Table 1) and seedling growth (Table 2). These results corroborate the hypothesis that with age, the allelopathic potential of sorghum and its hybrids decreases.

Lettuce, tomato, and pigweed seed germination and growth inhibition by sorghum-sudangrass extracts suggest that this hybrid may have the potential to model plant communities and serve as an important alien species when introduced in fragile natural ecosystems such as Cerrado (Bais et al., 2006). When introduced into native ecosystems, sorghum-sudangrass may play a role as an alien species with high biomass production potential and allelochemical production able to suppresses the growth of other species (Trezzi & Vidal, 2004; Vidal & Trezzi, 2004).



Table 2 - Lettuce, tomato, purslane, and pigweed plant length after being treated with aqueous extracts of sorghum-sudangrass root and shoot harvested at 5 growth stages §

Extract concentration (g L ⁻¹)	Sorghum-sudangrass age (days after emergence)				
	10	20	30	40	50
	Germination (%)				
	Lettuce				
0	95.00 Aa	95.00 Aa	94.58 Aa	95.83 Aa	94.17 Aa
0.4R*	95.42 Aa	94.17 Aa	87.08 Aab	95.83 Aa	95.83 Aa
2R	83.33 ABab	94.58 Aa	87.08 Aa	93.33 Aa	92.50 Aa
10R	82.92 ABab	92.08 Aa	83.75 ABab	95.00 Aa	95.83 Aa
0.4S	91.25 Aa	94.17 Aa	87.08 Aab	95.83 Aa	96.67 Aa
2S	82.92 ABab	95.00 Aa	88.75 Aa	93.33 Aa	95.83 Aa
10S	64.58 Cb	84.58 ABa	89.58 Aa	92.50 Aa	91.67 Aa
	Tomato				
0	66.67 Aa	76.25 Aa	72.50 Aa	70.00 Aa	66.67 Aa
0.4R	70.41 Aa	69.17 Aa	68.75 Aa	75.83 Aa	67.50 Aa
2R	75.00 Aa	71.67 Aa	66.67 Aa	73.33 Aa	72.50 Aa
10R	70.00 Aa	68.10 Aa	71.66 Aa	68.33 Aa	71.67 Aa
0.4S	65.00 Aa	71.25 Aa	65.42 Aa	75.83 Aa	67.50 Aa
2S	60.83 Aa	73.75 Aa	68.33 Aa	69.17 Aa	68.33 Aa
10S	12.08 Bb	3.75 Bc	65.00 Aa	71.67 Aa	66.67 Aa
	Purslane				
0	81.25 Aa	75.00 Aa	78.75 Aa	72.50 Aa	78.33 Aa
0.4R	79.17 Aa	80.83 Aa	78.33 ABa	82.50 Aa	76.67 Aa
2R	81.67 Aa	81.67 Aa	80.83 Aa	69.17 ABb	82.50 Aa
10R	79.17 Aa	80.83 Aa	78.33 ABa	80.00 Aa	77.50 Aa
0.4S	79.17 Aa	74.17 Aa	80.83 Aa	75.83 Aa	76.67 Aa
2S	80.00 Aa	74.58 Aab	89.17 Aa	78.33 Aab	80.83 Aa
10S	76.25 Aa	75.42 Aa	74.17 ABa	73.33 Aa	84.17 Aa
	Pigweed				
0	90.00 Aa	91.67 Aa	92.08 Aa	90.00 ABa	96.67 Aa
0.4R	89.58 Aa	88.33 Aa	90.42 Aa	91.67 ABa	89.17 ABa
2R	87.50 Aa	86.25 Aa	83.33 Aab	94.17 Aa	88.33 ABa
10R	89.05 Ab	87.50 Ab	88.33 Ab	98.33 Aa	90.00 Aa
0.4S	80.42 Aab	86.67 Aab	89.58 Aa	96.67 Aa	93.33 Aa
2S	82.08 Aab	85.42 Aab	89.58 Aab	97.50 Aa	95.83 Aa
10S	2.92 Bb	2.08 Bb	85.83 Aa	95.00 Aa	90.00 Aa

§ Different concentrations (0.4, 2, 10 g L⁻¹) of root (R) or shoot (S) extracts from sorghum-sudangrass plants harvested at 5 different stages (10, 20, 30, 40, 50 days after germination) were used to treat the seeds of four plant species (lettuce, tomato, purslane, and pigweed). For each species, plant length in the same column followed by the same upper case letter was not significantly different, while values in the same rows followed by the same lower case letter were not significantly different by the Tukey test *pd* 5%.

* R = root extracts; S = shoot extracts.

Table 3 - Lettuce, tomato, purslane, and pigweed root length after being treated with root and shoot aqueous extracts from sorghum-sudangrass harvested at 5 growth stages[§]

Extract concentration (g L ⁻¹)	Sorghum-sudangrass age (days after emergence)				
	10	20	30	40	50
	Radicle (mm)				
	Lettuce				
0	18.01 Ab	13.53 Ac	13.39 Bc	19.89 Ab	27.67 Aa
0.4R*	16.02 ABa	7.46 Bb	15.07 ABa	9.82 CDb	9.25 Cb
2R	13.04 BCb	12.60 Ab	18.49 Aa	17.28 ABa	17.57 Ba
10R	10.09 Cb	12.00 Aab	15.53 ABa	13.60 BCab	9.96 Cb
0.4S	12.89 BCc	15.93 Abc	12.64 Bc	18.89 Ab	25.49 Aa
2S	3.33 Dc	12.14 Ab	13.46 Bab	16.42 ABa	15.82 Bab
10S	2.43 Dc	3.53 Bc	12.64 Ba	8.58 Db	7.82 Cb
	Tomato				
0	35.00 ABb	56.45 Aa	48.66 Aab	35.22 ABb	37.83 Ab
0.4R	34.35 ABb	51.75 Aa	48.53 Aa	17.67 BCb	26.90 ABb
2R	50.39 Aab	49.41 Aab	60.85 Aa	38.71 Abc	24.39 ABc
10R	33.85 ABbc	49.96 Aab	58.07 Aa	28.32 Abc	26.39 ABc
0.4S	20.28 BCd	59.20 Aa	48.92 Aab	42.28 Abc	28.30 ABcd
2S	11.21 CDd	60.41 Aa	47.82 Aab	24.50 ABCcd	33.59 ABbc
10S	1.64 Dbc	0.00 Bc	48.82 Aa	9.68 Cbc	16.96 Bb
	Purslane				
0	19.10 Aa	20.00 Aa	20.96 Aa	23.96 ABa	21.00 ABa
0.4R	19.07 Aa	9.39 BCb	11.60 Bb	11.96 CDb	7.85 Db
2R	18.80 ABab	15.39 ABb	24.21 Aa	17.57 Cb	16.10 BCb
10R	12.51 BCbc	17.53 Ab	26.96 Aa	24.89 Aa	11.63 CDc
0.4S	7.62 CDc	21.50 Ab	23.39 Aab	27.35 Aa	23.28 Aab
2S	2.44 Dd	20.57 Aab	24.82 Aa	18.21 BCbc	13.57 CDc
10S	2.50 Dc	4.88 Cbc	22.28 Aa	7.89 Dbc	9.86 CDb
	Pigweed				
0	29.53 Aab	27.75 Ab	37.75 Aa	31.85 Aab	31.43 Aab
0.4R	24.66 Aa	9.53 BCb	13.07 Bb	8.64 CDb	5.89 Bb
2R	23.32 ABb	24.32 Ab	38.74 Aab	32.26 Aab	26.18 Ab
10R	13.00 BCc	18.64 ABbc	34.71 Aa	22.71 ABb	15.57 Bbc
0.4S	12.64 Cc	25.17 Ab	35.43 Aa	31.10 Aab	26.35 Aab
2S	5.25 CDc	26.17 Aa	30.14 Aa	15.85 BCb	14.67 Bbc
10S	0.00 Dc	0.00 Cc	33.07 Aa	5.10 Dbc	10.60 Bb

§ Different concentrations (0.4, 2, 10 g L⁻¹) of root (R) or shoot (S) extracts from sorghum-sudangrass plants harvested at 5 different stages (10, 20, 30, 40, 50 days after germination) were used to treat the seeds of four plant species (lettuce, tomato, purslane, and pigweed). For each species, plant length in the same column followed by the same upper case letter was not significantly different, while values in the same rows followed by the same lower case letter were not significantly different by the Tukey test $p \leq 5\%$.

* R = root extracts; S = shoot extracts.



Table 4 - Lettuce, tomato, purslane, and pigweed shoot length after treatment with root and shoot aqueous extracts from sorghum-sudangrass harvested at 5 growth stages[§]

Extract concentration (g L ⁻¹)	Sorghum-sudangrass age (days after emergence)				
	10	20	30	40	50
	Shoots (mm)				
	Lettuce				
0	15.67 Abc	11.78 Bc	14.10 Bbc	18.07 Cb	24.03 BCb
0.4R*	16.32 Ab	19.96 Aab	18.68 Aab	21.92 ABCa	22.53 BCa
2R	16.25 Abc	15.39 Bbc	12.68 Cc	18.92 Cab	21.96 BCab
10R	16.80 Abc	11.28 Bd	13.35 Bcd	20.96 BCa	19.96 Ca
0.4S	13.64 ABb	12.93 Bb	15.46 ABCb	21.00 BCa	22.93 BCa
2S	10.85 Bc	13.28 Bbc	17.35 ABb	24.28 ABa	26.39 ABa
10S	0.00 Cd	6.68 Cc	17.00 ABbC	25.99 Aa	28.92 Aa
	Tomato				
0	24.89 ABa	23.50 Ba	28.15 Aa	23.34 Aa	21.12 Ca
0.4R	24.91 ABc	41.50 Aa	34.78 Aab	20.53 Ac	33.60 Ab
2R	24.25 Bb	26.75 Bab	32.02 Aa	24.17 Ab	23.42 BCb
10R	32.32 Aa	24.95 Bb	32.96 Aa	23.83 Ab	26.35 AabBC
0.4S	21.50 Bb	28.08 Bab	31.03 Aa	24.85 Aab	23.86 BCab
2S	13.42 Cb	26.12 Ba	31.89 Aa	25.46 Aa	29.77 ABa
10S	0.00 Da	0.00 Ca	33.28 Aa	24.64 Ab	30.03 ABab
	Purslane				
0	20.71 Aa	15.46 BCb	18.54 Cab	20.85 Da	20.28 Ba
0.4R	22.96 Aa	20.96 Aa	22.6A BCa	22.46 CDa	23.99 ABa
2R	22.80 Aab	16.10 Bc	21.07 ABCb	25.92 ABa	22.71 Bab
10R	23.41 Aa	15.14 BCc	19.03 BCbc	22.39 CDab	23.75 Ba
0.4S	15.69 Bc	13.93 BCc	20.39 ABCb	24.53 ABCab	24.93 ABa
2S	10.69 Cd	15.46 BCc	23.28 ABb	28.53 Aa	23.28 Bb
10S	3.28 Dc	10.78 Cb	24.86 Aa	24.46 ABCa	28.64 Aa
	Pigweed				
0	20.00 Ba	15.50 Ba	18.25 Ca	19.46 Ca	18.67 Ca
0.4R	22.30 Bb	35.93 Aa	32.17 Aab	29.96 Bb	23.21 BCa
2R	23.42 Ba	17.85 Bb	19.50 BCab	23.11 Ca	19.43 Cab
10R	35.28 Aa	16.46 Bc	23.10 BCb	22.60 Cb	22.00 BCb
0.4S	23.03 Bab	14.64 Bc	20.61 BCab	23.85 Ca	18.53 Cbc
2S	19.10 Bc	18.25 Bc	24.36 Bb	29.57 Ba	25.82 Bab
10S	0.00 Cb	0.00 Cb	33.64 Aa	35.75 Aa	32.35 Aa

§ Different concentrations (0.4, 2, 10 g L⁻¹) of root (R) or shoot (S) extracts from sorghum-sudangrass plants harvested at 5 different stages (10, 20, 30, 40, 50 days after germination) were used to treat the seeds of four plant species (lettuce, tomato, purslane, and pigweed). For each species, plant length in the same column followed by the same upper case letter was not significantly different, while values in the same rows followed by the same lower case letter were not significantly different by the Tukey test $p < 5\%$.

* R = root extracts; S = shoot extracts.

LITERATURE CITED

- AN, M. et al. Mathematical modeling of allelopathy: II. The dynamics of allelochemicals from living plants in the environment. **Ecol. Modeling**, v. 116, n.1, p. 53-66, 2003.
- BAIS, H. P. et al. The role of root exudates in rhizosphere and other organisms. **An. Rev. Plant Biol.**, v. 57, p. 233-66, 2006
- BÀRBERI, P.; MAZZONCINI, M. Changes in weed community composition as influenced by cover crop and management system in continuous corn. **Weed Sci.**, v. 49, n. 4, p. 491-499, 2001.
- BARBOSA, T. M. L. et al. Caracterização química e efeitos alelopáticos de exsudatos radiculares de plântulas de sorgo sobre alface. **Planta Daninha**, v. 16, n. 2, p. 153-162, 1998.
- BEN-HAMMOUDA, M. et al. A chemical basis for differential allelopathic potential of sorghum hybrids on wheat. **J. Chem. Ecol.**, v. 21, n. 6, p. 775-786, 1995.
- BLYTH, C. R. Approximate binomial confidence limits. **J. Am. Stat.**, v. 81, n. 385, p. 843-855, 1996.
- CORREIA, N. M.; CENTURION, M. A. P. C.; ALVES, P. L. C. A. Influência de extratos aquosos de sorgo sobre a germinação e desenvolvimento de plântulas de soja. **Ci. Rural**, v. 35, n. 3, p. 498-503, 2005a.
- CORREIA, N. M.; SOUZA, I. F.; KLINK, U. P. Palha de sorgo associada ao herbicida imazamox no controle de plantas daninhas na cultura da soja em sucessão. **Planta Daninha**, v.23, n. 3, p. 483-489, 2005b.
- CZARNOTA, M. A. et al. Mode of action, localization of production, chemical nature, and activity of sorgoleone: a potent PSII inhibitor in *Sorghum* spp. root exudates. **Weed Technol.**, v. 15, n. 4, p. 813-825, 2001.
- DAYAN, F. E. Factors modulating the levels of the allelochemical sorgoleone in *Sorghum bicolor*. **Planta**, v. 224, n. 2, p. 339-346, 2006.
- EVERITT, J. H.; ALANIZ, M. A.; LEE, J. B. Seed germination characteristics of *Kochia scoparia*. **J. Range Manag.**, v. 36, n. 5, p. 646-648, 1983.
- FERREIRA, D. F. Análises estatísticas por meio do Sisvar para Windows versão 4.0. In: REUNIÃO ANUAL DA REGIÃO BRASILEIRA DA SOCIEDADE INTERNACIONAL DE BIOMETRIA, 45., São Carlos, 2000. **Proceedings...** São Carlos, Universidade Federal de São Carlos, 2000. p. 255-258.
- FORNEY, D. R.; FOY, C. L. Phytotoxicity of products from rhizospheres of a sorghum-sudangrass hybrid (*Sorghum bicolor* x *Sorghum sudanense*). **Weed Sci.**, v. 33, n. 5, p. 597-604, 1985.
- GONZALEZ, V. M. et al. Inhibition of a photosystem II electron transfer reaction by the natural product sorgoleone. **J. Agric. Food Chem.**, v. 45, n. 4, p.1415-1421, 1997.
- HEJL, A. M.; KOSTER, K. L. The allelochemical sorgoleone inhibits root H⁺-ATPase and water uptake. **J. Chem. Ecol.**, v. 30, n. 11, p. 2181-2191, 2004.
- LEEMIS, L. M.; TRIVEDI, K. S. A comparison of approximate interval estimators for the Bernoulli parameter. **Am. Stat.**, v. 50, n. 1, p. 63-68, 1996.
- MCGUIRE, A. **Cover crops for the Columbia basin**. Washington: Cooperative Extension Washington State University, 2003. 2 p. (EB1950E).
- MCKINNEY, D. E. et al. A preliminary study of dual use of cover crops: sorghum-sudangrass as both hay and summer cover crop for no-till organic cabbage. In: SOUTHERN CONSERVATION TILLAGE CONFERENCE FOR SUSTAINABLE AGRICULTURE, 26., 2004, Raleigh. **Proceedings...** Raleigh: 2004. 10p.
- MELLO, R. et al. Análise produtiva e qualitativa de um híbrido de sorgo interespecífico submetido a dois cortes. **R. Bras. Milho Sorgo**, v. 2, n. 1, p. 20-33, 2003.
- NETZLEY, D. H.; BUTLER, L. G. Roots of sorghum exude hydrophobic droplets containing biologically active components. **Crop Sci.**, v. 26, n. 4, p. 775-780, 1986.
- NICOLLIER, G. F.; POPE, D. F.; THOMPSON, A. C. Biological activity of dhurrin and other compounds from johnson grass (*Sorghum halepense*). **J. Agric. Food Chem.**, v. 31, n. 4, p. 748-751, 1983.
- OLIBONE, D. et al. Crescimento inicial da soja sob efeito de resíduos de sorgo. **Planta Daninha**, v. 24, n. 2, p. 255-261, 2006.
- PRATT, J. W. A normal approximation for binomial, F, beta, and other common, related tail probabilities, II. **J. Am. Statist.**, v. 63, n. 324, p. 1457-1483, 1968.
- RAUPP, A. A. A.; BRANÇÃO, N. Resultados de área de validação de tecnologia com sorgo de corte/pastejo submetido a pastejo rotativo 1999/2000. In: REUNIÃO TÉCNICA DO MILHO, 45.; REUNIÃO TÉCNICA DO SORGO, 28., 2000, Pelotas. **Anais...** Pelotas: Embrapa Clima Temperado, 2000. p. 673.



- ROTH, C. M.; SHROYER, P.; PAULSEN, G. M. Allelopathy of sorghum on wheat under several tillage systems. **Agron. J.**, v. 92, n. 5, p. 855-860, 2000.
- SENE, M.; GALLET, C.; DORÉ, T. Phenolic compounds in a Sahelian sorghum (*Sorghum bicolor*) genotype (ce(145-66)) and associated soils. **J. Chem. Ecol.**, v. 27, n. 1, p. 81-92, 2001.
- TOMICH, T. R. et al. Potencial forrageiro de híbridos de sorgo com capim-sudão. **Arq. Bras. Med. Veter. Zoot.**, v. 6, n. 2, p. 258-263, 2004.
- TREZZI, M. M.; VIDAL, R. A. Potencial de utilização de cobertura vegetal de sorgo e milho na supressão de plantas daninhas em condição de campo: II - efeitos da cobertura morta. **Planta Daninha**, v. 22, n. 1, 1-10, 2004.
- VIDAL, R. A.; TREZZI, M. M. Potencial da utilização de coberturas vegetais de sorgo e milho na supressão de plantas daninhas em condição de campo: I - Plantas em desenvolvimento vegetativo. **Planta Daninha**, v. 22, n. 2, p. 217-223, 2004.
- WESTON, L.A. Utilization of allelopathy for weed management in agroecosystems. **Agron. J.**, v. 88, p.860-866, 1996.
- WESTON, L.A.; HARMON, R.; MUELLER, S. Allelopathic potential of sorghum-sudangrass hybrid (sudex). **J. Chem. Ecol.**, v. 15, n. 6, p. 1855-1865, 1989.
- WOODHEAD, S. Environmental and biotic factors affecting the phenolic content of different cultivars of Sorghum bicolor. **J. Chem. Ecol.**, v. 7, n. 7, p. 1035-1047, 1981.
- XUAN, T. D. et al. Methods to determine allelopathic potential of crop plants for weed control. **Allelopathy J.**, v. 13, n. 2, p. 149-164, 2004.
- ZAGO, C. P. Utilização do sorgo na alimentação de ruminantes. In: EMPRESA BRASILEIRA DE PESQUISA AGROPECUÁRIA - Centro Nacional de Pesquisa de Milho e Sorgo. **Manejo cultural do sorgo para forragem**. Sete Lagoas: 1997. p. 9-26. (Embrapa-CNPMS. Circular Técnica, 17).

